

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
3 March 2005 (03.03.2005)

PCT

(10) International Publication Number
WO 2005/018541 A2

(51) International Patent Classification⁷: **A61K**

European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(21) International Application Number:
PCT/US2004/022059

(22) International Filing Date: 8 July 2004 (08.07.2004)

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/486,549 11 July 2003 (11.07.2003) US

(71) Applicant (for all designated States except US): **PHARMACIA CORPORATION** [US/US]; Global Patent Department, 700 Chesterfield Parkway West, Chesterfield, Missouri 63017-1732 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **STEPHENSON, Diane, T.** [US/US]; 1527 North Road, Groton, CT 06340 (US).

(74) Agents: **PETRILO, Kathleen, M.** et al.; Senniger Powers, #1 Metropolitan Square, 16th Floor, St. Louis, MO 63102 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

Published:

— without international search report and to be republished upon receipt of that report

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMPOSITIONS OF A CYCLOOXYGENASE-2 SELECTIVE INHIBITOR AND A SEROTONIN-MODULATING AGENT FOR THE TREATMENT OF CENTRAL NERVOUS SYSTEM DAMAGE

(57) Abstract: The present invention provides compositions and methods for the treatment of central nervous system damage in a subject. More particularly, the invention provides a combination therapy for the treatment of a central nervous system ischemic condition or a central nervous system traumatic injury comprising the administration to a subject of a serotonin-modulating agent in combination with a cyclooxygenase-2 selective inhibitor.

WO 2005/018541 A2

COMPOSITIONS OF A CYCLOOXYGENASE-2 SELECTIVE INHIBITOR AND A SEROTONIN-MODULATING AGENT FOR THE TREATMENT OF CENTRAL NERVOUS SYSTEM DAMAGE

FIELD OF THE INVENTION

[0001] The present invention provides compositions and methods for the treatment of central nervous system damage. More particularly, the invention is directed toward a combination therapy for the treatment or prevention of ischemic-mediated central nervous system damage including ischemic stroke, or central nervous system damage resulting from traumatic injury, comprising the administration to a subject of a serotonin-modulating agent in combination with a cyclooxygenase-2 selective inhibitor.

BACKGROUND OF THE INVENTION

[0002] The continued increase in the incidence of ischemic-mediated central nervous system damage, including ischemic stroke, provides compelling evidence that there is a continuing need for better treatment strategies. Stroke, for example, is consistently the second or the third leading cause of death annually and the leading producer of disability among adults in the United States and western countries. Moreover, roughly 10% of patients with stroke become heavily handicapped, often needing attendant care.

[0003] Within the 1990's decade, the pathology underlying ischemic-mediated central nervous system injury was elucidated. Generally speaking, the normal amount of perfusion to brain gray matter is 60 to 70 mL/100 g of brain tissue/min. Death of central nervous system cells typically occurs only when the flow of blood falls below a certain level (approximately 8-10 mL/100 g of brain tissue/min) while at slightly higher levels the tissue remains alive but not able to function. For example, most strokes culminate in a core area of cell death (infarction) in which blood flow is so drastically reduced that the cells usually cannot recover. This threshold seems to occur when cerebral blood flow is 20 percent of normal or less. Without neuroprotective agents, nerve cells facing 80 to 100 percent ischemia will be irreversibly damaged within a few minutes. Surrounding the ischemic core is another area of tissue called the "ischemic penumbra" or "transitional zone" in which cerebral blood flow is between 20 and 50 percent of normal. Cells in this area are endangered, but not yet irreversibly damaged. Thus in the acute stroke, the affected central core brain tissue may die while the more peripheral tissues remain alive for many years after the initial insult, depending on the amount of blood the brain tissue receives.

[0004] At the cellular level, if left untreated, brain or spinal cell injury and death progress in a stepwise manner rapidly within the core infarction, and over time within the ischemic penumbra. Without adequate blood supply, brain or spinal cells lose their ability to produce energy, particularly adenosine triphosphate (ATP). When this energy failure occurs, brain or spinal cells become damaged and will die if critical thresholds are reached. Immediate cell death within the ischemic core is typically necrotic, while cell death in the penumbra may be either necrotic or apoptotic. It is believed that there are an immense number of mechanisms at work causing brain or spinal cell damage and death following energy failure. Each of these mechanisms represents a potential route for intervention. One of the ways brain cells respond to energy failure is by elevating the concentration of intracellular calcium. Worsening this and driving the concentrations to dangerous levels is the process of excitotoxicity, in which brain cells release excessive amounts of glutamate, a neurotransmitter. This stimulates chemical and electrical activities in receptors on other brain cells, which leads to the degradation and destruction of vital cellular structures. Brain cells ultimately die as a result of the actions of calcium-activated proteases (enzymes which digest cell proteins), lipases (enzymes which digest cell membranes) and free radicals formed as a result of the ischemic cascade.


[0005] Interventions have been directed toward salvaging the ischemic penumbra and reducing its size. Restoration of blood flow is the first step toward rescuing the tissue within the penumbra. Therefore, timely recanalization of an occluded vessel to restore perfusion in both the penumbra and in the ischemic core is one treatment option employed. Partial recanalization also markedly reduces the size of the penumbra as well. Moreover, intravenous tissue plasminogen activator and other thrombolytic agents have been shown to have clinical benefit if they are administered within a few hours of symptom onset. Beyond this narrow time window, however, the likelihood of beneficial effects is reduced and hemorrhagic complications related to thrombolytic agents become excessive, seriously compromising their therapeutic value. Hypothermia decreases the size of the ischemic insult in both anecdotal clinical and laboratory reports. In addition, a wide variety of agents have been shown to reduce infarct volume in animal models. These agents include pharmacologic interventions that involve thrombolysis, calcium channel blockade, and cell membrane receptor antagonism. Successful treatment of stroke victims remains a high-unmet medical need. To date, no effective neuroprotective therapy exists to treat stroke. There is a continuing need for improved treatment regimes following ischemic-mediated central nervous system injury.

[0006] Neuroprotective agents have been shown to extend the time during which neurons within the ischemic penumbra remain viable (Albers, (1997) Am. J. Cardiol. 80(4C):4d-10d). Toward that end, several studies indicate that treatment with a serotonin-modulating agent following ischemic-mediated central nervous system injury may be beneficial. In particular, it has been suggested that several serotonin-modulating agents have shown neuroprotective effect in animal models of ischemia. In one study, for example, it was demonstrated that modulation of the serotonin system exerts a neuroprotective effect on delayed neuronal death after transient forebrain ischemia (Klisch, et al., (2003) Neurosci. Lett.; May 15(342)(1-2):25-28). Another study demonstrated a reduction in infarct volume to ischemic rabbits administered a serotonin-modulating agent (Shimizu, et al., (2002) J. Am. Coll. Cardiol.; Oct 2 40(7):1347-55). A further study suggests that serotonin-modulating agent administration is beneficial in traumatic brain injury (Kline, et al., (2002) Neurosci. Lett.; Nov 29(333)(3):179-82).

SUMMARY OF THE INVENTION

[0007] Among the several aspects of the invention is provided a method and a composition for the treatment of reduced blood flow to the central nervous system in a subject. The composition comprises a cyclooxygenase-2 selective inhibitor or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof and a serotonin-modulating agent or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof, and the method comprises administering to the subject a cyclooxygenase-2 selective inhibitor or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof in combination with a serotonin-modulating agent or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof.

[0008] In one embodiment, the cyclooxygenase-2 selective inhibitor is a member of the chromene class of compounds. For example, the chromene compound may be a compound of the formula:



[0009] wherein:

[0010] n is an integer which is 0, 1, 2, 3 or 4;

[0011] G is O, S or NR^a ;

[0012] R^a is alkyl;


[0013] R^1 is selected from the group consisting of H and aryl;

[0014] R^2 is selected from the group consisting of carboxyl, aminocarbonyl, alkylsulfonylaminocarbonyl and alkoxycarbonyl;

[0015] R^3 is selected from the group consisting of haloalkyl, alkyl, aralkyl, cycloalkyl and aryl optionally substituted with one or more radicals selected from alkylthio, nitro and alkylsulfonyl; and

[0016] each R^4 is independently selected from the group consisting of H, halo, alkyl, aralkyl, alkoxy, aryloxy, heteroaryloxy, aralkyloxy, heteroaralkyloxy, haloalkyl, haloalkoxy, alkylamino, arylamino, aralkylamino, heteroarylamino, heteroaralkylamino, nitro, amino, aminosulfonyl, alkylaminosulfonyl, arylaminosulfonyl, heteroarylaminosulfonyl, aralkylaminosulfonyl, heteroaralkylaminosulfonyl, heterocyclosulfonyl, alkylsulfonyl, hydroxyarylcarbonyl, nitroaryl, optionally substituted aryl, optionally substituted heteroaryl, aralkylcarbonyl, heteroarylcarbonyl, arylcarbonyl, aminocarbonyl, and alkylcarbonyl; or wherein R^4 together with the carbon atoms to which it is attached and the remainder of ring E forms a naphthyl radical.

[0017] In another embodiment, the cyclooxygenase-2 selective inhibitor or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof comprises a compound of the formula:



[0018] wherein:

[0019] A is selected from the group consisting of a partially unsaturated or unsaturated heterocyclyl ring and a partially unsaturated or unsaturated carbocyclic ring;

[0020] R^1 is selected from the group consisting of heterocyclyl, cycloalkyl, cycloalkenyl and aryl, wherein R^1 is optionally substituted at a substitutable position with one or more radicals selected from alkyl, haloalkyl, cyano, carboxyl,

alkoxycarbonyl, hydroxyl, hydroxyalkyl, haloalkoxy, amino, alkylamino, arylamino, nitro, alkoxyalkyl, alkylsulfinyl, halo, alkoxy and alkylthio;

[0021] R² is selected from the group consisting of methyl and amino; and

[0022] R³ is selected from the group consisting of H, halo, alkyl, alkenyl, alkynyl, oxo, cyano, carboxyl, cyanoalkyl, heterocycloxy, alkyloxy, alkylthio, alkylcarbonyl, cycloalkyl, aryl, haloalkyl, heterocyclyl, cycloalkenyl, aralkyl, heterocyclalkyl, acyl, alkylthioalkyl, hydroxyalkyl, alkoxycarbonyl, arylcarbonyl, aralkylcarbonyl, aralkenyl, alkoxyalkyl, arylthioalkyl, aryloxyalkyl, aralkylthioalkyl, aralkoxyalkyl, alkoxyaralkoxyalkyl, alkoxycarbonylalkyl, aminocarbonyl, aminocarbonylalkyl, alkylaminocarbonyl, N-arylaminocarbonyl, N-alkyl-N-arylaminocarbonyl, alkylaminocarbonylalkyl, carboxyalkyl, alkylamino, N-arylamino, N-aralkylamino, N-alkyl-N-aralkylamino, N-alkyl-N-arylamino, aminoalkyl, alkylaminoalkyl, N-arylaminalkyl, N-aralkylaminalkyl, N-alkyl-N-aralkylaminalkyl, N-alkyl-N-arylaminalkyl, aryloxy, aralkoxy, arylthio, aralkylthio, alkylsulfinyl, alkylsulfonyl, aminosulfonyl, alkylaminosulfonyl, N-arylaminosulfonyl, arylsulfonyl, and N-alkyl-N-arylaminosulfonyl.

[0023] In one embodiment, the serotonin-modulating agent is a serotonin receptor antagonist.

[0024] In another embodiment, the serotonin-modulating agent is a serotonin receptor agonist.

[0025] In another embodiment, the serotonin-modulating agent is a serotonin reuptake inhibitor.

[0026] Other aspects of the invention are described in more detail below.

ABBREVIATIONS AND DEFINITIONS

[0027] The term "acyl" is a radical provided by the residue after removal of hydroxyl from an organic acid. Examples of such acyl radicals include alkanoyl and aroyl radicals. Examples of such lower alkanoyl radicals include formyl, acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, pivaloyl, hexanoyl, and trifluoroacetyl.

[0028] The term "alkenyl" is a linear or branched radical having at least one carbon-carbon double bond of two to about twenty carbon atoms or, preferably, two to about twelve carbon atoms. More preferred alkyl radicals are "lower alkenyl" radicals having two to about six carbon atoms. Examples of alkenyl radicals include ethenyl, propenyl, allyl, propenyl, butenyl and 4-methylbutenyl. The terms "alkenyl" and "lower alkenyl" also are radicals having "cis" and "trans" orientations, or alternatively, "E" and "Z" orientations.

[0029] The term "cycloalkyl" is a saturated carbocyclic radical having three to twelve carbon atoms. More preferred cycloalkyl radicals are "lower cycloalkyl" radicals having three to about eight carbon atoms. Examples of such radicals include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

[0030] The terms "alkoxy" and "alkyloxy" are linear or branched oxy-containing radicals each having alkyl portions of one to about ten carbon atoms. More preferred alkoxy radicals are "lower alkoxy" radicals having one to six carbon atoms. Examples of such radicals include methoxy, ethoxy, propoxy, butoxy and tert-butoxy.

[0031] The term "alkoxyalkyl" is an alkyl radical having one or more alkoxy radicals attached to the alkyl radical, that is, to form monoalkoxyalkyl and dialkoxyalkyl radicals. The "alkoxy" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide haloalkoxy radicals. More preferred haloalkoxy radicals are "lower haloalkoxy" radicals having one to six carbon atoms and one or more halo radicals. Examples of such radicals include fluoromethoxy, chloromethoxy, trifluoromethoxy, trifluoroethoxy, fluoroethoxy and fluoropropoxy.

[0032] The term "alkoxycarbonyl" is a radical containing an alkoxy radical, as defined above, attached via an oxygen atom to a carbonyl radical. More preferred are "lower alkoxycarbonyl" radicals with alkyl portions having 1 to 6 carbons. Examples of such lower alkoxycarbonyl (ester) radicals include substituted or unsubstituted methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl and hexyloxycarbonyl.

[0033] Where used, either alone or within other terms such as "haloalkyl", "alkylsulfonyl", "alkoxyalkyl" and "hydroxyalkyl", the term "alkyl" is a linear, cyclic or branched radical having one to about twenty carbon atoms or, preferably, one to about twelve carbon atoms. More preferred alkyl radicals are "lower alkyl" radicals having one to about ten carbon atoms. Most preferred are lower alkyl radicals having one to about six carbon atoms. Examples of such radicals include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, iso-amyl, hexyl and the like.

[0034] The term "alkylamino" is an amino group that has been substituted with one or two alkyl radicals. Preferred are "lower N-alkylamino" radicals having alkyl portions having 1 to 6 carbon atoms. Suitable lower alkylamino may be mono or dialkylamino such as N-methylamino, N-ethylamino, N,N-dimethylamino, N,N-diethylamino or the like.

[0035] The term "alkylaminoalkyl" is a radical having one or more alkyl radicals attached to an aminoalkyl radical.

[0036] The term "alkylaminocarbonyl" is an aminocarbonyl group that has been substituted with one or two alkyl radicals on the amino nitrogen atom. Preferred are "N-

alkylaminocarbonyl" "N,N-dialkylaminocarbonyl" radicals. More preferred are "lower N-alkylaminocarbonyl" "lower N,N-dialkylaminocarbonyl" radicals with lower alkyl portions as defined above.

[0037] The terms "alkylcarbonyl", "arylcarbonyl" and "aralkylcarbonyl" include radicals having alkyl, aryl and aralkyl radicals, as defined above, attached to a carbonyl radical. Examples of such radicals include substituted or unsubstituted methylcarbonyl, ethylcarbonyl, phenylcarbonyl and benzylcarbonyl.

[0038] The term "alkylthio" is a radical containing a linear or branched alkyl radical, of one to about ten carbon atoms attached to a divalent sulfur atom. More preferred alkylthio radicals are "lower alkylthio" radicals having alkyl radicals of one to six carbon atoms. Examples of such lower alkylthio radicals are methylthio, ethylthio, propylthio, butylthio and hexylthio.

[0039] The term "alkylthioalkyl" is a radical containing an alkylthio radical attached through the divalent sulfur atom to an alkyl radical of one to about ten carbon atoms. More preferred alkylthioalkyl radicals are "lower alkylthioalkyl" radicals having alkyl radicals of one to six carbon atoms. Examples of such lower alkylthioalkyl radicals include methylthiomethyl.

[0040] The term "alkylsulfinyl" is a radical containing a linear or branched alkyl radical, of one to ten carbon atoms, attached to a divalent $-S(=O)-$ radical. More preferred alkylsulfinyl radicals are "lower alkylsulfinyl" radicals having alkyl radicals of one to six carbon atoms. Examples of such lower alkylsulfinyl radicals include methylsulfinyl, ethylsulfinyl, butylsulfinyl and hexylsulfinyl.

[0041] The term "alkynyl" is a linear or branched radical having two to about twenty carbon atoms or, preferably, two to about twelve carbon atoms. More preferred alkynyl radicals are "lower alkynyl" radicals having two to about ten carbon atoms. Most preferred are lower alkynyl radicals having two to about six carbon atoms. Examples of such radicals include propargyl, butynyl, and the like.

[0042] The term "aminoalkyl" is an alkyl radical substituted with one or more amino radicals. More preferred are "lower aminoalkyl" radicals. Examples of such radicals include aminomethyl, aminoethyl, and the like.

[0043] The term "aminocarbonyl" is an amide group of the formula $-C(=O)NH_2$.

[0044] The term "aralkoxy" is an aralkyl radical attached through an oxygen atom to other radicals.

[0045] The term "aralkoxyalkyl" is an aralkoxy radical attached through an oxygen atom to an alkyl radical.

[0046] The term "aralkyl" is an aryl-substituted alkyl radical such as benzyl, diphenylmethyl, triphenylmethyl, phenylethyl, and diphenylethyl. The aryl in said aralkyl may be additionally substituted with halo, alkyl, alkoxy, haloalkyl and haloalkoxy. The terms benzyl and phenylmethyl are interchangeable.

[0047] The term "aralkylamino" is an aralkyl radical attached through an amino nitrogen atom to other radicals. The terms "N-arylaminoalkyl" and "N-aryl-N-alkyl-aminoalkyl" are amino groups which have been substituted with one aryl radical or one aryl and one alkyl radical, respectively, and having the amino group attached to an alkyl radical. Examples of such radicals include N-phenylaminomethyl and N-phenyl-N-methylaminomethyl.

[0048] The term "aralkylthio" is an aralkyl radical attached to a sulfur atom.

[0049] The term "aralkylthioalkyl" is an aralkylthio radical attached through a sulfur atom to an alkyl radical.

[0050] The term "aroyl" is an aryl radical with a carbonyl radical as defined above. Examples of aroyl include benzoyl, naphthoyl, and the like and the aryl in said aroyl may be additionally substituted.

[0051] The term "aryl", alone or in combination, is a carbocyclic aromatic system containing one, two or three rings wherein such rings may be attached together in a pendent manner or may be fused. The term "aryl" includes aromatic radicals such as phenyl, naphthyl, tetrahydronaphthyl, indane and biphenyl. Aryl moieties may also be substituted at a substitutable position with one or more substituents selected independently from alkyl, alkoxyalkyl, alkylaminoalkyl, carboxyalkyl, alkoxycarbonylalkyl, aminocarbonylalkyl, alkoxy, aralkoxy, hydroxyl, amino, halo, nitro, alkylamino, acyl, cyano, carboxy, aminocarbonyl, alkoxycarbonyl and aralkoxycarbonyl.

[0052] The term "arylamino" is an amino group, which has been substituted with one or two aryl radicals, such as N-phenylamino. The "arylamino" radicals may be further substituted on the aryl ring portion of the radical.

[0053] The term "aryloxyalkyl" is a radical having an aryl radical attached to an alkyl radical through a divalent oxygen atom.

[0054] The term "arylthioalkyl" is a radical having an aryl radical attached to an alkyl radical through a divalent sulfur atom.

[0055] The term "carbonyl", whether used alone or with other terms, such as "alkoxycarbonyl", is $-(C=O)-$.

[0056] The terms "carboxy" or "carboxyl", whether used alone or with other terms, such as "carboxyalkyl", is $-CO_2H$.

[0057] The term "carboxyalkyl" is an alkyl radical substituted with a carboxy radical. More preferred are "lower carboxyalkyl" which are lower alkyl radicals as defined above, and may be additionally substituted on the alkyl radical with halo. Examples of such lower carboxyalkyl radicals include carboxymethyl, carboxyethyl and carboxypropyl.

[0058] The term "cycloalkenyl" is a partially unsaturated carbocyclic radical having three to twelve carbon atoms. More preferred cycloalkenyl radicals are "lower cycloalkenyl" radicals having four to about eight carbon atoms. Examples of such radicals include cyclobutenyl, cyclopentenyl, cyclopentadienyl, and cyclohexenyl.

[0059] The term "cyclooxygenase-2 selective inhibitor" is a compound able to selectively inhibit cyclooxygenase-2 over cyclooxygenase-1. Typically, it includes compounds that have a cyclooxygenase-2 IC_{50} of less than about 0.2 micro molar, and also have a selectivity ratio of cyclooxygenase-1 (COX-1) IC_{50} to cyclooxygenase-2 (COX-2) IC_{50} of at least about 5, more typically of at least about 50, and even more typically, of at least about 100. Moreover, the cyclooxygenase-2 selective inhibitors as described herein have a cyclooxygenase-1 IC_{50} of greater than about 1 micro molar, and more preferably of greater than 10 micro molar. Inhibitors of the cyclooxygenase pathway in the metabolism of arachidonic acid used in the present method may inhibit enzyme activity through a variety of mechanisms. By the way of example, and without limitation, the inhibitors used in the methods described herein may block the enzyme activity directly by acting as a substrate for the enzyme.

[0060] The term "halo" is a halogen such as fluorine, chlorine, bromine or iodine.

[0061] The term "haloalkyl" is a radical wherein any one or more of the alkyl carbon atoms is substituted with halo as defined above. Specifically included are monohaloalkyl, dihaloalkyl and polyhaloalkyl radicals. A monohaloalkyl radical, for one example, may have either an iodo, bromo, chloro or fluoro atom within the radical. Dihalogen and polyhaloalkyl radicals may have two or more of the same halo atoms or a combination of different halo radicals. "Lower haloalkyl" is a radical having 1-6 carbon atoms. Examples of haloalkyl radicals include fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluorochloromethyl, dichlorofluoromethyl, difluoroethyl, difluoropropyl, dichloroethyl and dichloropropyl.

[0062] The term "heteroaryl" is an unsaturated heterocyclyl radical. Examples of unsaturated heterocyclyl radicals, also termed "heteroaryl" radicals include unsaturated 3 to 6 membered heteromonocyclic group containing 1 to 4 nitrogen atoms, for example, pyrrolyl, pyrrolinyl, imidazolyl, pyrazolyl, pyridyl, pyrimidyl, pyrazinyl, pyridazinyl, triazolyl

(e.g., 4H-1,2,4-triazolyl, 1H-1,2,3-triazolyl, 2H-1,2,3-triazolyl, etc.) tetrazolyl (e.g. 1H-tetrazolyl, 2H-tetrazolyl, etc.), etc.; unsaturated condensed heterocyclyl group containing 1 to 5 nitrogen atoms, for example, indolyl, isoindolyl, indolizynyl, benzimidazolyl, quinolyl, isoquinolyl, indazolyl, benzotriazolyl, tetrazolopyridazinyl (e.g., tetrazolo[1,5-b]pyridazinyl, etc.), etc.; unsaturated 3 to 6-membered heteromonocyclic group containing an oxygen atom, for example, pyranyl, furyl, etc.; unsaturated 3 to 6-membered heteromonocyclic group containing a sulfur atom, for example, thienyl, etc.; unsaturated 3- to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms, for example, oxazolyl, isoxazolyl, oxadiazolyl (e.g., 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,5-oxadiazolyl, etc.) etc.; unsaturated condensed heterocyclyl group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms (e.g. benzoxazolyl, benzoxadiazolyl, etc.); unsaturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms, for example, thiazolyl, thiadiazolyl (e.g., 1,2,4- thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, etc.) etc.; unsaturated condensed heterocyclyl group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms (e.g., benzothiazolyl, benzothiadiazolyl, etc.) and the like. The term also includes radicals where heterocyclyl radicals are fused with aryl radicals. Examples of such fused bicyclic radicals include benzofuran, benzothiophene, and the like. Said "heterocyclyl group" may have 1 to 3 substituents such as alkyl, hydroxyl, halo, alkoxy, oxo, amino and alkylamino.

[0063] The term "heterocyclyl" is a saturated, partially unsaturated and unsaturated heteroatom-containing ring-shaped radical, where the heteroatoms may be selected from nitrogen, sulfur and oxygen. Examples of saturated heterocyclyl radicals include saturated 3 to 6-membered heteromonocyclic group containing 1 to 4 nitrogen atoms (e.g. pyrrolidinyl, imidazolidinyl, piperidino, piperazinyl, etc.); saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 oxygen atoms and 1 to 3 nitrogen atoms (e.g. morpholiny, etc.); saturated 3 to 6-membered heteromonocyclic group containing 1 to 2 sulfur atoms and 1 to 3 nitrogen atoms (e.g., thiazolidinyl, etc.). Examples of partially unsaturated heterocyclyl radicals include dihydrothiophene, dihydropyran, dihydrofuran and dihydrothiazole.

[0064] The term "heterocyclylalkyl" is a saturated and partially unsaturated heterocyclyl-substituted alkyl radical, such as pyrrolidinylmethyl, and heteroaryl-substituted alkyl radicals, such as pyridylmethyl, quinolylmethyl, thienylmethyl, furylethyl, and quinolylethyl. The heteroaryl in said heteroalkyl may be additionally substituted with halo, alkyl, alkoxy, haloalkyl and haloalkoxy.

[0065] The term "hydrido" is a single hydrogen atom (H). This hydrido radical may be attached, for example, to an oxygen atom to form a hydroxyl radical or two hydrido radicals may be attached to a carbon atom to form a methylene (-CH₂-) radical.

[0066] The term "hydroxyalkyl" is a linear or branched alkyl radical having one to about ten carbon atoms any one of which may be substituted with one or more hydroxyl radicals. More preferred hydroxyalkyl radicals are "lower hydroxyalkyl" radicals having one to six carbon atoms and one or more hydroxyl radicals. Examples of such radicals include hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl and hydroxyhexyl.

[0067] The term "pharmaceutically acceptable" is used adjectivally herein to mean that the modified noun is appropriate for use in a pharmaceutical product; that is the "pharmaceutically acceptable" material is relatively safe and/or non-toxic, though not necessarily providing a separable therapeutic benefit by itself. Pharmaceutically acceptable cations include metallic ions and organic ions. More preferred metallic ions include, but are not limited to appropriate alkali metal salts, alkaline earth metal salts and other physiologically acceptable metal ions. Exemplary ions include aluminum, calcium, lithium, magnesium, potassium, sodium and zinc in their usual valences. Preferred organic ions include protonated tertiary amines and quaternary ammonium cations, including in part, trimethylamine, diethylamine, N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. Exemplary pharmaceutically acceptable acids include without limitation hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, methanesulfonic acid, acetic acid, formic acid, tartaric acid, maleic acid, malic acid, citric acid, isocitric acid, succinic acid, lactic acid, gluconic acid, glucuronic acid, pyruvic acid, oxalacetic acid, fumaric acid, propionic acid, aspartic acid, glutamic acid, benzoic acid, and the like.

[0068] The term "prodrug" refers to a chemical compound that can be converted into a therapeutic compound by metabolic or simple chemical processes within the body of the subject. For example, a class of prodrugs of COX-2 inhibitors is described in US Patent No. 5,932,598, herein incorporated by reference.

[0069] The term "serotonin-modulating agent," unless otherwise indicated herein, includes a number of suitable agents that alter serotonin levels by interacting with serotonin receptor sites (5HT₁-5HT₇) in the body. The term also includes a number of suitable agents that either alter the biological activity of serotonin or result in a change in the amount of biologically active serotonin. Modulation can be an increase or decrease in

activity, a change in binding characteristics, or any other change in the biological, functional, or immunological properties of biologically active serotonin.

[0070] The term "subject" for purposes of treatment includes any human or animal who has reduced blood flow to the central nervous system. The subject can be a domestic livestock species, a laboratory animal species, a zoo animal or a companion animal. In one embodiment, the subject is a mammal. In another embodiment, the mammal is a human being.

[0071] The term "sulfonyl", whether used alone or linked to other terms such as alkylsulfonyl, is a divalent radical $-\text{SO}_2-$. "Alkylsulfonyl" is an alkyl radical attached to a sulfonyl radical, where alkyl is defined as above. More preferred alkylsulfonyl radicals are "lower alkylsulfonyl" radicals having one to six carbon atoms. Examples of such lower alkylsulfonyl radicals include methylsulfonyl, ethylsulfonyl and propylsulfonyl. The "alkylsulfonyl" radicals may be further substituted with one or more halo atoms, such as fluoro, chloro or bromo, to provide haloalkylsulfonyl radicals. The terms "sulfamyl", "aminosulfonyl" and "sulfonamidyl" are $\text{NH}_2\text{O}_2\text{S}-$.

[0072] The phrase "therapeutically-effective" is intended to qualify the amount of each agent (i.e. the amount of cyclooxygenase-2 selective inhibitor and the amount of serotonin-modulating agent) which will achieve the goal of improvement in disorder severity and the frequency of incidence over no treatment or treatment of each agent by itself.

[0073] The term "thrombotic event" or "thromboembolic event" includes, but is not limited to arterial thrombosis, including stent and graft thrombosis, cardiac thrombosis, coronary thrombosis, heart valve thrombosis, pulmonary thrombosis and venous thrombosis. Cardiac thrombosis is thrombosis in the heart. Pulmonary thrombosis is thrombosis in the lung. Arterial thrombosis is thrombosis in an artery. Coronary thrombosis is the development of an obstructive thrombus in a coronary artery, often causing sudden death or a myocardial infarction. Venous thrombosis is thrombosis in a vein. Heart valve thrombosis is a thrombosis on a heart valve. Stent thrombosis is thrombosis resulting from and/or located in the vicinity of a vascular stent. Graft thrombosis is thrombosis resulting from and/or located in the vicinity of an implanted graft, particularly a vascular graft. A thrombotic event as used herein is meant to embrace both a local thrombotic event and a distal thrombotic event occurring anywhere within the body (e.g., a thromboembolic event such as for example an embolic stroke).

[0074] The term "treat" or "treatment" as used herein, includes administration of the combination therapy to a subject known to have central nervous system damage. In

other aspects, it also includes either preventing the onset of clinically evident central nervous system damage altogether or preventing the onset of preclinically evident stage of central nervous system damage. This definition includes prophylactic treatment.


[0075] The term "vaso-occlusive event" includes a partial occlusion (including a narrowing) or complete occlusion of a blood vessel, a stent or a vascular graft. A vaso-occlusive event intends to embrace thrombotic or thromboembolic events, and the vascular occlusion disorders or conditions to which they give rise. Thus, a vaso-occlusive event is intended to embrace all vascular occlusive disorders resulting in partial or total vessel occlusion from thrombotic or thromboembolic events.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0076] The present invention provides a combination therapy comprising the administration to a subject of a therapeutically effective amount of a COX-2 selective inhibitor in combination with a therapeutically effective amount of a serotonin-modulating agent. The combination therapy is used to treat or prevent damage to a central nervous system cell resulting from a reduction in blood flow or traumatic injury. When administered as part of a combination therapy, the COX-2 selective inhibitor together with the serotonin-modulating agent provide enhanced treatment options as compared to administration of either the serotonin-modulating agent or the COX-2 selective inhibitor alone.


CYCLOOXYGENASE-2 SELECTIVE INHIBITORS

[0077] A number of suitable cyclooxygenase-2 selective inhibitors or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof, may be employed in the composition of the current invention. In one embodiment, the cyclooxygenase-2 selective inhibitor can be, for example, the cyclooxygenase-2 selective inhibitor meloxicam, Formula B-1 (CAS registry number 71125-38-7) or an isomer, a pharmaceutically acceptable salt, ester, or prodrug of a compound having Formula B-1.



[0078] In yet another embodiment, the cyclooxygenase-2 selective inhibitor is the cyclooxygenase-2 selective inhibitor, 6-[[5-(4-chlorobenzoyl)-1,4-dimethyl-1H-pyrrol-2-


yl]methyl]-3(2H)-pyridazinone, Formula B-2 (CAS registry number 179382-91-3) or an isomer, a pharmaceutically acceptable salt, ester, or prodrug of a compound having Formula B-2.



[0079] In still another embodiment the cyclooxygenase-2 selective inhibitor is a chromene compound that is a substituted benzopyran or a substituted benzopyran analog, and even more typically, selected from the group consisting of substituted benzothiopyrans, dihydroquinolines, dihydronaphthalenes or a compound having

[0080] Formula I shown below and possessing, by way of example and not limitation, the structures disclosed in Table 1. Furthermore, benzopyran cyclooxygenase-2 selective inhibitors useful in the practice of the present methods are described in U.S. Patent No. 6,034,256 and 6,077,850 herein incorporated by reference in their entirety.

[0081] In another embodiment, the cyclooxygenase-2 selective inhibitor is a chromene compound represented by Formula I or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof:



[0082] wherein:

[0083] n is an integer which is 0, 1, 2, 3 or 4;

[0084] G is O, S or NR^a;

[0085] R^a is alkyl;

[0086] R¹ is selected from the group consisting of H and aryl;

[0087] R² is selected from the group consisting of carboxyl, lower alkyl, lower aralkyl, aminocarbonyl, alkylsulfonylaminocarbonyl and alkoxycarbonyl;

[0088] R³ is selected from the group consisting of haloalkyl, alkyl, aralkyl, cycloalkyl and aryl optionally substituted with one or more radicals selected from the group consisting of alkylthio, nitro and alkylsulfonyl; and

[0089] each R^4 is independently selected from the group consisting of H, halo, alkyl, aralkyl, alkoxy, aryloxy, heteroaryloxy, aralkyloxy, heteroaralkyloxy, haloalkyl, haloalkoxy, alkylamino, arylamino, aralkylamino, heteroaryl amino, heteroarylalkylamino, nitro, amino, aminosulfonyl, alkylaminosulfonyl, arylaminosulfonyl, heteroarylaminosulfonyl, aralkylaminosulfonyl, heteroaralkylaminosulfonyl, heterocyclosulfonyl, alkylsulfonyl, hydroxyarylcarbonyl, nitroaryl, optionally substituted aryl, optionally substituted heteroaryl, aralkylcarbonyl, heteroarylcarbonyl, arylcarbonyl, aminocarbonyl, and alkylcarbonyl; or R^4 together with the carbon atoms to which it is attached and the remainder of ring E forms a naphthyl radical.

[0090] The cyclooxygenase-2 selective inhibitor may also be a compound of Formula (I) or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof,

[0091] wherein:

[0092] n is an integer which is 0, 1, 2, 3 or 4;

[0093] G is O, S or NR^a ;

[0094] R^a is alkyl;

[0095] R^1 is H;

[0096] R^2 is selected from the group consisting of carboxyl, aminocarbonyl, alkylsulfonylaminocarbonyl and alkoxy carbonyl;

[0097] R^3 is selected from the group consisting of haloalkyl, alkyl, aralkyl, cycloalkyl and aryl optionally substituted with one or more radicals selected from the group consisting of alkylthio, nitro and alkylsulfonyl; and

[0098] each R^4 is independently selected from the group consisting of hydrido, halo, alkyl, aralkyl, alkoxy, aryloxy, heteroaryloxy, aralkyloxy, heteroaralkyloxy, haloalkyl, haloalkoxy, alkylamino, arylamino, aralkylamino, heteroaryl amino, heteroarylalkylamino, nitro, amino, aminosulfonyl, alkylaminosulfonyl, arylaminosulfonyl, heteroarylaminosulfonyl, aralkylaminosulfonyl, heteroaralkylaminosulfonyl, heterocyclosulfonyl, alkylsulfonyl, optionally substituted aryl, optionally substituted heteroaryl, aralkylcarbonyl, heteroarylcarbonyl, arylcarbonyl, aminocarbonyl, and alkylcarbonyl; or wherein R^4 together with the carbon atoms to which it is attached and the remainder of ring E forms a naphthyl radical.

[0099] In a further embodiment, the cyclooxygenase-2 selective inhibitor may also be a compound of Formula (I), or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof,

[0100] wherein:

[0101] n is an integer which is 0, 1, 2, 3 or 4;

[0102] G is oxygen or sulfur;
[0103] R¹ is H;
[0104] R² is carboxyl, lower alkyl, lower aralkyl or lower alkoxy carbonyl;
[0105] R³ is lower haloalkyl, lower cycloalkyl or phenyl; and
[0106] each R⁴ is independently H, halo, lower alkyl, lower alkoxy, lower haloalkyl, lower haloalkoxy, lower alkylamino, nitro, amino, aminosulfonyl, lower alkylaminosulfonyl, 5-membered heteroarylalkylaminosulfonyl, 6-membered heteroarylalkylaminosulfonyl, lower aralkylaminosulfonyl, 5-membered nitrogen-containing heterocyclosulfonyl, 6-membered-nitrogen containing heterocyclosulfonyl, lower alkylsulfonyl, optionally substituted phenyl, lower aralkylcarbonyl, or lower alkylcarbonyl; or R⁴ together with the carbon atoms to which it is attached and the remainder of ring E forms a naphthyl radical.

[0107] The cyclooxygenase-2 selective inhibitor may also be a compound of Formula (I) or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof,

[0108] wherein:
[0109] n is an integer which is 0, 1, 2, 3 or 4;
[0110] G is oxygen or sulfur;
[0111] R¹ is H;
[0112] R² is carboxyl;
[0113] R³ is lower haloalkyl; and
[0114] each R⁴ is independently H, halo, lower alkyl, lower haloalkyl, lower haloalkoxy, lower alkylamino, amino, aminosulfonyl, lower alkylaminosulfonyl, 5-membered heteroarylalkylaminosulfonyl, 6-membered heteroarylalkylaminosulfonyl, lower aralkylaminosulfonyl, lower alkylsulfonyl, 6-membered nitrogen-containing heterocyclosulfonyl, optionally substituted phenyl, lower aralkylcarbonyl, or lower alkylcarbonyl; or wherein R⁴ together with the carbon atoms to which it is attached and the remainder of ring E forms a naphthyl radical.

[0115] The cyclooxygenase-2 selective inhibitor may also be a compound of Formula (I) or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof,

[0116] wherein:
[0117] n is an integer which is 0, 1, 2, 3 or 4;
[0118] G is oxygen or sulfur;
[0119] R¹ is H;
[0120] R² is carboxyl;

[0121] R^3 is fluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, pentafluoroethyl, heptafluoropropyl, difluoroethyl, difluoropropyl, dichloroethyl, dichloropropyl, difluoromethyl, or trifluoromethyl; and

[0122] each R^4 is independently H, chloro, fluoro, bromo, iodo, methyl, ethyl, isopropyl, *tert*-butyl, butyl, isobutyl, pentyl, hexyl, methoxy, ethoxy, isopropoxy, *tert*butyloxy, trifluoromethyl, difluoromethyl, trifluoromethoxy, amino, N,N-dimethylamino, N,N-diethylamino, N-phenylmethylaminosulfonyl, N-phenylethylaminosulfonyl, N-(2-furylmethyl)aminosulfonyl, nitro, N,N-dimethylaminosulfonyl, aminosulfonyl, N-methylaminosulfonyl, N-ethylsulfonyl, 2,2-dimethylethylaminosulfonyl, N,N-dimethylaminosulfonyl, N-(2-methylpropyl)aminosulfonyl, N-morpholinosulfonyl, methylsulfonyl, benzylcarbonyl, 2,2-dimethylpropylcarbonyl, phenylacetyl or phenyl; or wherein R^4 together with the carbon atoms to which it is attached and the remainder of ring E forms a naphthyl radical.

[0123] The cyclooxygenase-2 selective inhibitor may also be a compound of Formula (I) or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof,

[0124] wherein:

[0125] n is an integer which is 0, 1, 2, 3 or 4;

[0126] G is oxygen or sulfur;

[0127] R^1 is H;

[0128] R^2 is carboxyl;

[0129] R^3 is trifluoromethyl or pentafluoroethyl; and

[0130] each R^4 is independently H, chloro, fluoro, bromo, iodo, methyl, ethyl, isopropyl, *tert*-butyl, methoxy, trifluoromethyl, trifluoromethoxy, N-phenylmethylaminosulfonyl, N-phenylethylaminosulfonyl, N-(2-furylmethyl)aminosulfonyl, N,N-dimethylaminosulfonyl, N-methylaminosulfonyl, N-(2,2-dimethylethyl)aminosulfonyl, dimethylaminosulfonyl, 2-methylpropylaminosulfonyl, N-morpholinosulfonyl, methylsulfonyl, benzylcarbonyl, or phenyl; or wherein R^4 together with the carbon atoms to which it is attached and the remainder of ring E forms a naphthyl radical.

[0131] In yet another embodiment, the cyclooxygenase-2 selective inhibitor used in connection with the method(s) of the present invention can also be a compound having the structure of Formula (I) or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof,

[0132] wherein:

[0133] n is 4;

[0134] G is O or S;

[0135] R^1 is H;

[0136] R^2 is CO_2H ;

[0137] R^3 is lower haloalkyl;


[0138] a first R^4 corresponding to R^9 is hydrido or halo;

[0139] a second R^4 corresponding to R^{10} is H, halo, lower alkyl, lower haloalkoxy, lower alkoxy, lower aralkylcarbonyl, lower dialkylaminosulfonyl, lower alkylaminosulfonyl, lower aralkylaminosulfonyl, lower heteroaralkylaminosulfonyl, 5-membered nitrogen-containing heterocyclosulfonyl, or 6-membered nitrogen-containing heterocyclosulfonyl;

[0140] a third R^4 corresponding to R^{11} is H, lower alkyl, halo, lower alkoxy, or aryl; and

[0141] a fourth R^4 corresponding to R^{12} is H, halo, lower alkyl, lower alkoxy, or aryl;

[0142] wherein Formula (I) is represented by Formula (Ia):



[0143] The cyclooxygenase-2 selective inhibitor used in connection with the method(s) of the present invention can also be a compound of having the structure of Formula (Ia) or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof,

[0144] wherein:

[0145] G is O or S;

[0146] R^3 is trifluoromethyl or pentafluoroethyl;

[0147] R^9 is H, chloro, or fluoro;





[0148] R^{10} is H, chloro, bromo, fluoro, iodo, methyl, tert-butyl, trifluoromethoxy, methoxy, benzylcarbonyl, dimethylaminosulfonyl, isopropylaminosulfonyl, methylaminosulfonyl, benzylaminosulfonyl, phenylethylaminosulfonyl, methylpropylaminosulfonyl, methylsulfonyl, or morpholinosulfonyl;






[0149] R^{11} is H, methyl, ethyl, isopropyl, tert-butyl, chloro, methoxy, diethylamino, or phenyl; and







[0150] R^{12} is H, chloro, bromo, fluoro, methyl, ethyl, tert-butyl, methoxy, or phenyl.

[0151] Examples of exemplary chromene cyclooxygenase-2 selective inhibitors are depicted in Table 1 below.


TABLE 1
EXAMPLES OF CHROMENE CYCLOOXYGENASE-2 SELECTIVE INHIBITORS AS EMBODIMENTS

<u>Compound Number</u>	<u>Structural Formula</u>
B-3	 <p>6-Nitro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid</p>
B-4	 <p>6-Chloro-8-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid</p>
B-5	 <p>((S)-6-Chloro-7-(1,1-dimethylethyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid</p>
B-6	 <p>2-Trifluoromethyl-2H-naphtho[2,3-b]pyran-3-carboxylic acid</p>

<u>Compound Number</u>	<u>Structural Formula</u>
B-7	 <p>6-Chloro-7-(4-nitrophenoxy)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid</p>
B-8	 <p>((S)-6,8-Dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid</p>
B-9	 <p>6-Chloro-2-(trifluoromethyl)-4-phenyl-2H-1-benzopyran-3-carboxylic acid</p>
B-10	 <p>6-(4-Hydroxybenzoyl)-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid</p>
B-11	 <p>2-(Trifluoromethyl)-6-[(trifluoromethyl)thio]-2H-1-benzothiopyran-3-carboxylic acid</p>

<u>Compound Number</u>	<u>Structural Formula</u>
B-12	 <p>6,8-Dichloro-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid</p>
B-13	 <p>6-(1,1-Dimethylethyl)-2-(trifluoromethyl)-2H-1-benzothiopyran-3-carboxylic acid</p>
B-14	 <p>6,7-Difluoro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid</p>
B-15	 <p>6-Chloro-1,2-dihydro-1-methyl-2-(trifluoromethyl)-3-quinolinecarboxylic acid</p>
B-16	 <p>6-Chloro-2-(trifluoromethyl)-1,2-dihydro[1,8]naphthyridine-3-carboxylic acid</p>
B-17	 <p>((S)-6-Chloro-1,2-dihydro-2-(trifluoromethyl)-3-quinolinecarboxylic acid</p>

[0152] In a further embodiment, the cyclooxygenase-2 selective inhibitor is selected from the class of tricyclic cyclooxygenase-2 selective inhibitors represented by the general structure of Formula II or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof,



[0153] wherein:

[0154] A is selected from the group consisting of a partially unsaturated or unsaturated heterocyclyl ring and a partially unsaturated or unsaturated carbocyclic ring;

[0155] R¹ is selected from the group consisting of heterocyclyl, cycloalkyl, cycloalkenyl and aryl, wherein R¹ is optionally substituted at a substitutable position with one or more radicals selected from alkyl, haloalkyl, cyano, carboxyl, alkoxy carbonyl, hydroxyl, hydroxyalkyl, haloalkoxy, amino, alkylamino, arylamino, nitro, alkoxyalkyl, alkylsulfinyl, halo, alkoxy and alkylthio;

[0156] R² is selected from the group consisting of methyl and amino; and



[0157] R³ is selected from the group consisting of H, halo, alkyl, alkenyl, alkynyl, oxo, cyano, carboxyl, cyanoalkyl, heterocycloxy, alkyloxy, alkylthio, alkylcarbonyl, cycloalkyl, aryl, haloalkyl, heterocyclyl, cycloalkenyl, aralkyl, heterocyclalkyl, acyl, alkylthioalkyl, hydroxyalkyl, alkoxy carbonyl, arylcarbonyl, aralkylcarbonyl, aralkenyl, alkoxyalkyl, arylthioalkyl, aryloxyalkyl, aralkylthioalkyl, aralkoxyalkyl, alkoxyaralkoxyalkyl, alkoxy carbonylalkyl, aminocarbonyl, aminocarbonylalkyl, alkylaminocarbonyl, N- arylaminocarbonyl, N-alkyl-N- arylaminocarbonyl, alkylaminocarbonylalkyl, carboxyalkyl, alkylamino, N-aryl amino, N-aralkylamino, N-alkyl-N-aralkylamino, N-alkyl-N-aryl amino, aminoalkyl, alkylaminoalkyl, N-aryl aminoalkyl, N-aralkylaminoalkyl, N-alkyl-N-aralkylaminoalkyl, N-alkyl-N-aryl aminoalkyl, aryloxy, aralkoxy, arylthio, aralkylthio, alkylsulfinyl, alkylsulfonyl, aminosulfonyl, alkylaminosulfonyl, N-arylaminosulfonyl, arylsulfonyl, and N-alkyl-N-arylaminosulfonyl.

[0158] In another embodiment, the cyclooxygenase-2 selective inhibitor represented by the above Formula II is selected from the group of compounds illustrated in

Table 2, consisting of celecoxib (B-18; U.S. Patent No. 5,466,823; CAS No. 169590-42-5), valdecoxib (B-19; U.S. Patent No. 5,633,272; CAS No. 181695-72-7), deracoxib (B-20; U.S. Patent No. 5,521,207; CAS No. 169590-41-4), rofecoxib (B-21; CAS No. 162011-90-7), etoricoxib (MK-663; B-22; PCT publication WO 98/03484), tilmacoxib (JTE-522; B-23; CAS No. 180200-68-4), and cimicoxib (UR-8880; B23a; CAS No. 265114-23-6).


TABLE 2
EXAMPLES OF TRICYCLIC CYCLOOXYGENASE-2 SELECTIVE INHIBITORS AS EMBODIMENTS

<u>Compound Number</u>	<u>Structural Formula</u>
B-18	
B-19	
B-20	
B-21	
B-22	

<u>Compound Number</u>	<u>Structural Formula</u>
B-23	
B-23a	

[0159] In still another embodiment, the cyclooxygenase-2 selective inhibitor is selected from the group consisting of celecoxib, rofecoxib and etoricoxib.


[0160] In yet another embodiment, the cyclooxygenase-2 selective inhibitor is parecoxib (B-24, U.S. Patent No. 5,932,598, CAS No. 198470-84-7), which is a therapeutically effective prodrug of the tricyclic cyclooxygenase-2 selective inhibitor valdecoxib, B-19, may be advantageously employed as a source of a cyclooxygenase inhibitor (US 5,932,598, herein incorporated by reference).



B-24


[0161] One form of parecoxib is sodium parecoxib.

[0162] In another embodiment of the invention, the compound having the formula B-25 or an isomer, a pharmaceutically acceptable salt, ester, or prodrug of a compound having formula B-25 that has been previously described in International Publication number WO 00/24719 (which is herein incorporated by reference) is another tricyclic cyclooxygenase-2 selective inhibitor that may be advantageously employed.




B-25

[0163] Another cyclooxygenase-2 selective inhibitor that is useful in connection with the method(s) of the present invention is N-(2-cyclohexyloxynitrophenyl)-methane sulfonamide (NS-398) having a structure shown below as B-26, or an isomer, a pharmaceutically acceptable salt, ester, or prodrug of a compound having formula B-26.



B-26

[0164] In yet a further embodiment, the cyclooxygenase-2 selective inhibitor used in connection with the method(s) of the present invention can be selected from the class of phenylacetic acid derivative cyclooxygenase-2 selective inhibitors represented by the general structure of Formula (III) or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof:



(III)

[0165] wherein:

[0166] R^{16} is methyl or ethyl;

[0167] R^{17} is chloro or fluoro;

[0168] R^{18} is hydrogen or fluoro;

[0169] R^{19} is hydrogen, fluoro, chloro, methyl, ethyl, methoxy, ethoxy or hydroxy;

[0170] R^{20} is hydrogen or fluoro; and

[0171] R^{21} is chloro, fluoro, trifluoromethyl or methyl, provided, however, that each of R^{17} , R^{18} , R^{20} and R^{21} is not fluoro when R^{16} is ethyl and R^{19} is H.

[0172] Another phenylacetic acid derivative cyclooxygenase-2 selective inhibitor used in connection with the method(s) of the present invention is a compound that has the designation of COX 189 (lumiracoxib; B-211) and that has the structure shown in Formula (III) or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof wherein:


[0173] R^{16} is ethyl;

[0174] R^{17} and R^{19} are chloro;

[0175] R^{18} and R^{20} are hydrogen; and

[0176] R^{21} is methyl.

[0177] In yet another embodiment, the cyclooxygenase-2 selective inhibitor is represented by Formula (IV) or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof:



[0178] wherein:

[0179] X is O or S;

[0180] J is a carbocycle or a heterocycle;


[0181] R^{22} is NHSO_2CH_3 or F;

[0182] R^{23} is H, NO_2 , or F; and

[0183] R^{24} is H, NHSO_2CH_3 , or $(\text{SO}_2\text{CH}_3)\text{C}_6\text{H}_4$.

[0184] According to another embodiment, the cyclooxygenase-2 selective inhibitors used in the present method(s) have the structural Formula (V) or an isomer, a

pharmaceutically acceptable salt, ester, or prodrug thereof:



[0185] wherein:

[0186] T and M are independently phenyl, naphthyl, a radical derived from a heterocycle comprising 5 to 6 members and possessing from 1 to 4 heteroatoms, or a radical derived from a saturated hydrocarbon ring having from 3 to 7 carbon atoms;

[0187] R^{25} , R^{26} , R^{27} , and R^{28} are independently hydrogen, halogen, lower alkyl radical having from 1 to 6 carbon atoms, lower haloalkyl radical having from 1 to 6 carbon atoms, or an aromatic radical selected from the group consisting of phenyl, naphthyl, thienyl, furyl and pyridyl; or

[0188] R^{25} and R^{26} , together with the carbon atom to which they are attached, form a carbonyl or a saturated hydrocarbon ring having from 3 to 7 carbon atoms; or

[0189] R^{27} and R^{28} , together with the carbon atom to which they are attached, form a carbonyl or a saturated hydrocarbon ring having from 3 to 7 carbon atoms;

[0190] Q^1 , Q^2 , L^1 or L^2 are independently hydrogen, halogen, lower alkyl having from 1 to 6 carbon atoms, trifluoromethyl, lower methoxy having from 1 to 6 carbon atoms, alkylsulfinyl or alkylsulfonyl; and

[0191] at least one of Q^1 , Q^2 , L^1 or L^2 is in the para position and is $-S(O)_n-R$, wherein n is 0, 1, or 2 and R is a lower alkyl radical having 1 to 6 carbon atoms or a lower haloalkyl radical having from 1 to 6 carbon atoms, or an $-SO_2NH_2$; or Q^1 and Q^2 together form methylenedioxy; or L^1 and L^2 together form methylenedioxy.

[0192] In another embodiment, the compounds N-(2-cyclohexyloxynitrophenyl) methane sulfonamide, and (E)-4-[(4-methylphenyl)(tetrahydro-2-oxo-3-furanylidene) methyl]benzenesulfonamide or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof having the structure of Formula (V) are employed as cyclooxygenase-2 selective inhibitors.

[0193] In a further embodiment, compounds that are useful for the cyclooxygenase-2 selective inhibitor or an isomer, a pharmaceutically acceptable salt,

ester, or prodrug thereof used in connection with the method(s) of the present invention, the structures for which are set forth in Table 3 below, include, but are not limited to:

- [0194] 6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-27);
- [0195] 6-chloro-7-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-28);
- [0196] 8-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-29);
- [0197] 6-chloro-8-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-30);
- [0198] 2-trifluoromethyl-3H-naphtho[2,1-b]pyran-3-carboxylic acid (B-31);
- [0199] 7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-32);
- [0200] 6-bromo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-33);
- [0201] 8-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-34);
- [0202] 6-trifluoromethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-35);
- [0203] 5,7-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-36);
- [0204] 8-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-37);
- [0205] 7,8-dimethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-38);
- [0206] 6,8-bis(dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-39);
- [0207] 7-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-40);
- [0208] 7-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-41);
- [0209] 6-chloro-7-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-42);
- [0210] 6-chloro-8-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-43);
- [0211] 6-chloro-7-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-44);
- [0212] 6,7-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-45);
- [0213] 6,8-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-46);
- [0214] 6-chloro-8-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-47);

- [0215] 8-chloro-6-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-48)
- [0216] 8-chloro-6-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-49);
- [0217] 6-bromo-8-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-50);
- [0218] 8-bromo-6-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-51);
- [0219] 8-bromo-6-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-52);
- [0220] 8-bromo-5-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-53);
- [0221] 6-chloro-8-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-54);
- [0222] 6-bromo-8-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-55);
- [0223] 6-[[[(phenylmethyl)amino]sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-56);
- [0224] 6-[(dimethylamino)sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-57);
- [0225] 6-[(methylamino)sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-58);
- [0226] 6-[(4-morpholino)sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-59);
- [0227] 6-[(1,1-dimethylethyl)aminosulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-60);
- [0228] 6-[(2-methylpropyl)aminosulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-61);
- [0229] 6-methylsulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-62);
- [0230] 8-chloro-6-[[[(phenylmethyl)amino]sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-63);
- [0231] 6-phenylacetyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-64);
- [0232] 6,8-dibromo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-65);

[0233] 8-chloro-5,6-dimethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-66);

[0234] 6,8-dichloro-(*S*)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-67);

[0235] 6-benzylsulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-68);

[0236] 6-[[*N*-(2-furylmethyl)amino]sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-69);

[0237] 6-[[*N*-(2-phenylethyl)amino]sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-70);

[0238] 6-iodo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-71);

[0239] 7-(1,1-dimethylethyl)-2-pentafluoroethyl-2H-1-benzopyran-3-carboxylic acid (B-72);

[0240] 6-chloro-2-trifluoromethyl-2H-1-benzothiopyran-3-carboxylic acid (B-73);

[0241] 3-[(3-chloro-phenyl)-(4-methanesulfonyl-phenyl)-methylene]-dihydro-furan-2-one or BMS-347070 (B-74);

[0242] 8-acetyl-3-(4-fluorophenyl)-2-(4-methylsulfonyl)phenyl-imidazo(1,2-*a*)pyridine (B-75);

[0243] 5,5-dimethyl-4-(4-methylsulfonyl)phenyl-3-phenyl-2-(5H)-furanone (B-76);

[0244] 5-(4-fluorophenyl)-1-[4-(methylsulfonyl)phenyl]-3-(trifluoromethyl)pyrazole (B-77);

[0245] 4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]-1-phenyl-3-(trifluoromethyl)pyrazole (B-78);

[0246] 4-(5-(4-chlorophenyl)-3-(4-methoxyphenyl)-1H-pyrazol-1-yl) benzenesulfonamide (B-79);

[0247] 4-(3,5-bis(4-methylphenyl)-1H-pyrazol-1-yl) benzenesulfonamide (B-80);

[0248] 4-(5-(4-chlorophenyl)-3-phenyl-1H-pyrazol-1-yl) benzenesulfonamide (B-81);

[0249] 4-(3,5-bis(4-methoxyphenyl)-1H-pyrazol-1-yl) benzenesulfonamide (B-82);

[0250] 4-(5-(4-chlorophenyl)-3-(4-methylphenyl)-1H-pyrazol-1-yl) benzenesulfonamide (B-83);

[0251] 4-(5-(4-chlorophenyl)-3-(4-nitrophenyl)-1H-pyrazol-1-yl) benzenesulfonamide (B-84);

- [0252] 4-(5-(4-chlorophenyl)-3-(5-chloro-2-thienyl)-1H-pyrazol-1-yl) benzenesulfonamide (B-85);
- [0253] 4-(4-chloro-3,5-diphenyl-1H-pyrazol-1-yl)benzenesulfonamide (B-86);
- [0254] 4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide (B-87);
- [0255] 4-[5-phenyl-3-(trifluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide (B-88);
- [0256] 4-[5-(4-fluorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide (B-89);
- [0257] 4-[5-(4-methoxyphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide (B-90);
- [0258] 4-[5-(4-chlorophenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide (B-91);
- [0259] 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide (B-92);
- [0260] 4-[4-chloro-5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide (B-93);
- [0261] 4-[3-(difluoromethyl)-5-(4-methylphenyl)-1H-pyrazol-1-yl] benzenesulfonamide (B-94);
- [0262] 4-[3-(difluoromethyl)-5-phenyl-1H-pyrazol-1-yl] benzenesulfonamide (B-95);
- [0263] 4-[3-(difluoromethyl)-5-(4-methoxyphenyl)-1H-pyrazol-1-yl] benzenesulfonamide (B-96);
- [0264] 4-[3-cyano-5-(4-fluorophenyl)-1H-pyrazol-1-yl] benzenesulfonamide (B-97);
- [0265] 4-[3-(difluoromethyl)-5-(3-fluoro-4-methoxyphenyl)-1H-pyrazol-1-yl] benzenesulfonamide (B-98);
- [0266] 4-[5-(3-fluoro-4-methoxyphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide (B-99);
- [0267] 4-[4-chloro-5-phenyl-1H-pyrazol-1-yl]benzenesulfonamide (B-100);
- [0268] 4-[5-(4-chlorophenyl)-3-(hydroxymethyl)-1H-pyrazol-1-yl] benzenesulfonamide (B-101);
- [0269] 4-[5-(4-(N,N-dimethylamino)phenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide (B-102);
- [0270] 5-(4-fluorophenyl)-6-[4-(methylsulfonyl)phenyl]spiro[2.4]hept-

- [0271] 5-ene (B-103);
- [0272] 4-[6-(4-fluorophenyl)spiro[2.4]hept-5-en-5-yl] benzenesulfonamide (B-104);
- [0273] 6-(4-fluorophenyl)-7-[4-(methylsulfonyl)phenyl]spiro[3.4]oct-
- [0274] 6-ene (B-105);
- [0275] 5-(3-chloro-4-methoxyphenyl)-6-[4-(methylsulfonyl) phenyl]spiro [2.4]hept-5-ene (B-106);
- [0276] 4-[6-(3-chloro-4-methoxyphenyl)spiro[2.4]hept-5-en-5-yl] benzenesulfonamide (B-107);
- [0277] 5-(3,5-dichloro-4-methoxyphenyl)-6-[4-(methylsulfonyl)phenyl]spiro [2.4]hept-5-ene (B-108);
- [0278] 5-(3-chloro-4-fluorophenyl)-6-[4-(methylsulfonyl)phenyl] spiro[2.4]hept-5-ene (B-109);
- [0279] 4-[6-(3,4-dichlorophenyl)spiro[2.4]hept-5-en-5-yl] benzenesulfonamide (B-110);
- [0280] 2-(3-chloro-4-fluorophenyl)-4-(4-fluorophenyl)-5-(4-methylsulfonyl phenyl)thiazole (B-111);
- [0281] 2-(2-chlorophenyl)-4-(4-fluorophenyl)-5-(4-methylsulfonyl phenyl)thiazole (B-112);
- [0282] 5-(4-fluorophenyl)-4-(4-methylsulfonylphenyl)-2-methylthiazole (B-113);
- [0283] 4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-2-trifluoromethylthiazole (B-114);
- [0284] 4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-2-(2-thienyl)thiazole (B-115);
- [0285] 4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-2-benzylaminothiazole (B-116);
- [0286] 4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-2-(1-propylamino) thiazole (B-117);
- [0287] 2-[(3,5-dichlorophenoxy)methyl]-4-(4-fluorophenyl)-5-[4-(methyl sulfonyl)phenyl]thiazole (B-118);
- [0288] 5-(4-fluorophenyl)-4-(4-methylsulfonylphenyl)-2-trifluoromethylthiazole (B-119);
- [0289] 1-methylsulfonyl-4-[1,1-dimethyl-4-(4-fluorophenyl)cyclopenta-2,4-dien-3-yl]benzene (B-120);

- [0290] 4-[4-(4-fluorophenyl)-1,1-dimethylcyclopenta-2,4-dien-3-yl] benzenesulfonamide (B-121);
- [0291] 5-(4-fluorophenyl)-6-[4-(methylsulfonyl)phenyl]spiro[2.4]hepta-4,6-diene (B-122);
- [0292] 4-[6-(4-fluorophenyl)spiro[2.4]hepta-4,6-dien-5-yl] benzenesulfonamide (B-123);
- [0293] 6-(4-fluorophenyl)-2-methoxy-5-[4-(methylsulfonyl)phenyl]-pyridine-3-carbonitrile (B-124);
- [0294] 2-bromo-6-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]-pyridine-3-carbonitrile (B-125);
- [0295] 6-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]-2-phenyl-pyridine-3-carbonitrile (B-126);
- [0296] 4-[2-(4-methylpyridin-2-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl] benzenesulfonamide (B-127);
- [0297] 4-[2-(5-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl] benzenesulfonamide (B-128);
- [0298] 4-[2-(2-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl] benzenesulfonamide (B-129);
- [0299] 3-[1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazol-2-yl] pyridine (B-130);
- [0300] 2-[1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazol-
- [0301] 2-yl]pyridine (B-131);
- [0302] 2-methyl-4-[1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazol-2-yl]pyridine (B-132);
- [0303] 2-methyl-6-[1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazol-2-yl]pyridine (B-133);
- [0304] 4-[2-(6-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl] benzenesulfonamide (B-134);
- [0305] 2-(3,4-difluorophenyl)-1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazole (B-135);
- [0306] 4-[2-(4-methylphenyl)-4-(trifluoromethyl)-1H-imidazol-1-yl] benzenesulfonamide (B-136);
- [0307] 2-(4-chlorophenyl)-1-[4-(methylsulfonyl)phenyl]-4-methyl-1H-imidazole (B-137);

[0308] 2-(4-chlorophenyl)-1-[4-(methylsulfonyl)phenyl]-4-phenyl-1H-imidazole (B-138);

[0309] 2-(4-chlorophenyl)-4-(4-fluorophenyl)-1-[4-(methylsulfonyl)phenyl]-1H-imidazole (B-139);

[0310] 2-(3-fluoro-4-methoxyphenyl)-1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazole (B-140);

[0311] 1-[4-(methylsulfonyl)phenyl]-2-phenyl-4-trifluoromethyl-1H-imidazole (B-141);

[0312] 2-(4-methylphenyl)-1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazole (B-142);

[0313] 4-[2-(3-chloro-4-methylphenyl)-4-(trifluoromethyl)-1H-imidazol-1-yl] benzenesulfonamide (B-143);

[0314] 2-(3-fluoro-5-methylphenyl)-1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazole (B-144);

[0315] 4-[2-(3-fluoro-5-methylphenyl)-4-(trifluoromethyl)-1H-imidazol-1-yl] benzenesulfonamide (B-145);

[0316] 2-(3-methylphenyl)-1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazole (B-146);

[0317] 4-[2-(3-methylphenyl)-4-trifluoromethyl-1H-imidazol-1-yl] benzene sulfonamide (B-147);

[0318] 1-[4-(methylsulfonyl)phenyl]-2-(3-chlorophenyl)-4-trifluoromethyl-1H-imidazole (B-148);

[0319] 4-[2-(3-chlorophenyl)-4-trifluoromethyl-1H-imidazol-1-yl] benzenesulfonamide (B-149);

[0320] 4-[2-phenyl-4-trifluoromethyl-1H-imidazol-1-yl] benzenesulfonamide (B-150);

[0321] 4-[2-(4-methoxy-3-chlorophenyl)-4-trifluoromethyl-1H-imidazol-1-yl] benzenesulfonamide (B-151);

[0322] 1-allyl-4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)-1H-pyrazole (B-152);

[0323] 4-[1-ethyl-4-(4-fluorophenyl)-5-(trifluoromethyl)-1H-pyrazol-3-yl] benzenesulfonamide (B-153);

[0324] N-phenyl-[4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)-1H-pyrazol-1-yl]acetamide (B-154);

[0325] ethyl [4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)-1H-pyrazol-1-yl]acetate (B-155);

[0326] 4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-1-(2-phenylethyl)-1H-pyrazole (B-156);

[0327] 4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-1-(2-phenylethyl)-5-(trifluoromethyl)pyrazole (B-157);

[0328] 1-ethyl-4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)-1H-pyrazole (B-158);

[0329] 5-(4-fluorophenyl)-4-(4-methylsulfonylphenyl)-2-trifluoromethyl-1H-imidazole (B-159);

[0330] 4-[4-(methylsulfonyl)phenyl]-5-(2-thiophenyl)-2-(trifluoromethyl)-1H-imidazole (B-160);

[0331] 5-(4-fluorophenyl)-2-methoxy-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyridine (B-161);

[0332] 2-ethoxy-5-(4-fluorophenyl)-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyridine (B-162);

[0333] 5-(4-fluorophenyl)-4-[4-(methylsulfonyl)phenyl]-2-(2-propynyloxy)-6-(trifluoromethyl)pyridine (B-163);

[0334] 2-bromo-5-(4-fluorophenyl)-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyridine (B-164);

[0335] 4-[2-(3-chloro-4-methoxyphenyl)-4,5-difluorophenyl]benzenesulfonamide (B-165);

[0336] 1-(4-fluorophenyl)-2-[4-(methylsulfonyl)phenyl]benzene (B-166);

[0337] 5-difluoromethyl-4-(4-methylsulfonylphenyl)-3-phenylisoxazole (B-167);

[0338] 4-[3-ethyl-5-phenylisoxazol-4-yl]benzenesulfonamide (B-168);

[0339] 4-[5-difluoromethyl-3-phenylisoxazol-4-yl]benzenesulfonamide (B-169);

[0340] 4-[5-hydroxymethyl-3-phenylisoxazol-4-yl]benzenesulfonamide (B-170);

[0341] 4-[5-methyl-3-phenyl-isoxazol-4-yl]benzenesulfonamide (B-171);

[0342] 1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl) benzene (B-172);

[0343] 1-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl) benzene (B-173);

[0344] 1-[2-(4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl) benzene (B-174);

[0345] 1-[2-(2,4-dichlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl) benzene (B-175);

[0346] 1-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl) benzene (B-176);

[0347] 1-[2-(4-methylthiophenyl)cyclopenten-1-yl]-4-(methyl sulfonyl)benzene (B-177);

[0348] 1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl) benzene (B-178);

[0349] 4-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]benzene sulfonamide (B-179);

[0350] 1-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl) benzene (B-180);

[0351] 4-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]benzene sulfonamide (B-181);

[0352] 4-[2-(4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide (B-182);

[0353] 4-[2-(4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide (B-183);

[0354] 1-[2-(4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl) benzene (B-184);

[0355] 1-[2-(2,3-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl) benzene (B-185);

[0356] 4-[2-(3-fluoro-4-methoxyphenyl)cyclopenten-1-yl] benzenesulfonamide (B-186);

[0357] 1-[2-(3-chloro-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl) benzene (B-187);

[0358] 4-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl] benzenesulfonamide (B-188);

[0359] 4-[2-(2-methylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide (B-189);

[0360] ethyl 2-[4-(4-fluorophenyl)-5-[4-(methylsulfonyl) phenyl]oxazol-2-yl]-2-benzyl-acetate (B-190);

[0361] 2-[4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]oxazol-2-yl] acetic acid (B-191);

[0362] 2-(*tert*-butyl)-4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl] oxazole (B-192);

[0363] 4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]-2-phenyloxazole (B-193);

[0364] 4-(4-fluorophenyl)-2-methyl-5-[4-(methylsulfonyl)phenyl]oxazole (B-194);

- [0365] 4-[5-(3-fluoro-4-methoxyphenyl)-2-trifluoromethyl-4-oxazolyl] benzenesulfonamide (B-195);
- [0366] 6-chloro-7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-196);
- [0367] 6-chloro-8-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid (B-197);
- [0368] 5,5-dimethyl-3-(3-fluorophenyl)-4-methylsulfonyl-2(5H)-furanone (B-198);
- [0369] 6-chloro-2-trifluoromethyl-2H-1-benzothiopyran-3-carboxylic acid (B-199);
- [0370] 4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl] benzene sulfonamide (B-200);
- [0371] 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl] benzene sulfonamide (B-201);
- [0372] 4-[5-(3-fluoro-4-methoxyphenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide (B-202);
- [0373] 3-[1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazol-2-yl]pyridine (B-203);
- [0374] 2-methyl-5-[1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazol-2-yl]pyridine (B-204);
- [0375] 4-[2-(5-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl] benzenesulfonamide (B-205);
- [0376] 4-[5-methyl-3-phenylisoxazol-4-yl]benzenesulfonamide (B-206);
- [0377] 4-[5-hydroxymethyl-3-phenylisoxazol-4-yl]benzenesulfonamide (B-207);
- [0378] [2-trifluoromethyl-5-(3,4-difluorophenyl)-4-oxazolyl] benzenesulfonamide (B-208);
- [0379] 4-[2-methyl-4-phenyl-5-oxazolyl]benzenesulfonamide (B-209);
- [0380] 4-[5-(2-fluoro-4-methoxyphenyl)-2-trifluoromethyl-4-oxazolyl] benzenesulfonamide (B-210);
- [0381] [2-(2-chloro-6-fluoro-phenylamino)-5-methyl-phenyl]-acetic acid or COX 189 (lumiracoxib; B-211);
- [0382] N-(4-Nitro-2-phenoxy-phenyl)-methanesulfonamide or nimesulide (B-212);
- [0383] N-[6-(2,4-difluoro-phenoxy)-1-oxo-indan-5-yl]-methanesulfonamide or flosulide (B-213);
- [0384] N-[6-(2,4-Difluoro-phenylsulfanyl)-1-oxo-1H-inden-5-yl]-methanesulfonamide, sodium salt (B-214);




- [0385] N-[5-(4-fluoro-phenylsulfanyl)-thiophen-2-yl]-methanesulfonamide (B-215);
- [0386] 3-(3,4-Difluoro-phenoxy)-4-(4-methanesulfonyl-phenyl)-5-methyl-5-(2,2,2-trifluoro-ethyl)-5H-furan-2-one (B-216);
- [0387] (5Z)-2-amino-5-[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl] methylene]-4(5H)-thiazolone (B-217);
- [0388] CS-502 (B-218);
- [0389] LAS-34475 (B-219);
- [0390] LAS-34555 (B-220);
- [0391] S-33516 (B-221);
- [0392] SD-8381 (B-222);
- [0393] L-783003 (B-223);
- [0394] N-[3-(formylamino)-4-oxo-6-phenoxy-4H-1-benzopyran-7-yl]-methanesulfonamide (B-224);
- [0395] D-1367 (B-225);
- [0396] L-748731 (B-226);
- [0397] (6aR,10aR)-3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-6H-dibenzo[b,d]pyran-9-carboxylic acid (B-227);
- [0398] CGP-28238 (B-228);
- [0399] 4-[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene] dihydro-2-methyl-2H-1,2-oxazin-3(4H)-one or BF-389 (B-229);
- [0400] GR-253035 (B-230);
- [0401] 6-dioxo-9H-purin-8-yl-cinnamic acid (B-231);
- [0402] S-2474 (B-232);
- [0403] 4-[4-(methyl)-sulfonyl]phenyl]-3-phenyl-2(5H)-furanone;
- [0404] 4-(5-methyl-3-phenyl-4-isoxazolyl);
- [0405] 2-(6-methylpyrid-3-yl)-3-(4-methylsulfonylphenyl)-5-chloropyridine;
- [0406] 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl];
- [0407] N-[[4-(5-methyl-3-phenyl-4-isoxazolyl)phenyl]sulfonyl];
- [0408] 4-[5-(3-fluoro-4-methoxyphenyl)-3-difluoromethyl]-1H-pyrazol-
- [0409] 1-yl]benzenesulfonamide;
- [0410] (S)-6,8-dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid;
- [0411] 2-(3,4-difluorophenyl)-4-(3-hydroxy-3-methylbutoxy)-5-[4-(methylsulfonyl)phenyl]-3(2H)-pyridazinone;
- [0412] 2-trifluoromethyl-3H-naphtho[2,1-b]pyran-3-carboxylic acid;





[0413] 6-chloro-7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;





[0414] [2-(2,4-dichloro-6-ethyl-3,5-dimethyl-phenylamino)-5-propyl-phenyl]-acetic acid.





TABLE 3





EXAMPLES OF CYCLOOXYGENASE-2 SELECTIVE INHIBITORS AS EMBODIMENTS






<u>Compound Number</u>	<u>Structural Formula</u>
B-26	 <p>N-(2-cyclohexyloxynitrophenyl) methane sulfonamide or NS-398;</p>
B-27	 <p>6-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-28	 <p>6-chloro-7-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>





<u>Compound Number</u>	<u>Structural Formula</u>
B-29	 <p>8-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-30	 <p>6-chloro-8-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-31	 <p>2-trifluoromethyl-3H-naphtho[2,1-b]pyran-3-carboxylic acid;</p>
B-32	 <p>7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>





<u>Compound Number</u>	<u>Structural Formula</u>
B-33	 <p>6-bromo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-34	 <p>8-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-35	 <p>6-trifluoromethoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-36	 <p>5,7-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>





<u>Compound Number</u>	<u>Structural Formula</u>
B-37	 <p>8-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-38	 <p>7,8-dimethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-39	 <p>6,8-bis(dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-40	 <p>7-(1-methylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>





<u>Compound Number</u>	<u>Structural Formula</u>
B-41	 <p>7-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-42	 <p>6-chloro-7-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-43	 <p>6-chloro-8-ethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-44	 <p>6-chloro-7-phenyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>






<u>Compound Number</u>	<u>Structural Formula</u>
B-45	 <p>6,7-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-46	 <p>6,8-dichloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-47	 <p>6-chloro-8-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-48	 <p>8-chloro-6-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-49	 <p>8-chloro-6-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-50	 <p>6-bromo-8-chloro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-51	 <p>8-bromo-6-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-52	 <p>8-bromo-6-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-53	 <p>8-bromo-5-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-54	 <p>6-chloro-8-fluoro-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-55	 <p>6-bromo-8-methoxy-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-56	 <p>6-[[[(phenylmethyl)amino]sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-57	 <p>6-[(dimethylamino)sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-58	 <p>6-[(methylamino)sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-59	 <p>6-[(4-morpholino)sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-60	 <p>6-[(1,1-dimethylethyl)aminosulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-61	 <p>6-[(2-methylpropyl)aminosulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>



<u>Compound Number</u>	<u>Structural Formula</u>
B-62	 <p>6-methylsulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-63	 <p>8-chloro-6-[(phenylmethyl)amino]sulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-64	 <p>6-phenylacetyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-65	 <p>6,8-dibromo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-66	 <p>8-chloro-5,6-dimethyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-67	 <p>6,8-dichloro-(S)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-68	 <p>6-benzylsulfonyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-69	 <p>6-[[N-(2-furylmethyl)amino]sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-70	 <p>6-[[N-(2-phenylethyl)amino]sulfonyl]-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-71	 <p>6-iodo-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-72	 <p>7-(1,1-dimethylethyl)-2-pentafluoroethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-73	 <p>6-chloro-2-trifluoromethyl-2H-1-benzothiopyran-3-carboxylic acid;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-74	 <p>3-[(3-chloro-phenyl)-(4-methanesulfonyl-phenyl)-methylene]-dihydro-furan-2-one or BMS-347070;</p>
B-75	 <p>8-acetyl-3-(4-fluorophenyl)-2-(4-methylsulfonyl)phenyl-imidazo(1,2-a)pyridine;</p>
B-76	 <p>5,5-dimethyl-4-(4-methylsulfonyl)phenyl-3-phenyl-2-(5H)-furanone;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-77	 <p>5-(4-fluorophenyl)-1-[4-(methylsulfonyl)phenyl]-3-(trifluoromethyl)pyrazole;</p>
B-78	 <p>4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]-1-phenyl-3-(trifluoromethyl)pyrazole;</p>
B-79	 <p>4-(5-(4-chlorophenyl)-3-(4-methoxyphenyl)-1H-pyrazol-1-yl)benzenesulfonamide;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-80	 <p>4-(3,5-bis(4-methylphenyl)-1H-pyrazol-1-yl)benzenesulfonamide;</p>
B-81	 <p>4-(5-(4-chlorophenyl)-3-phenyl-1H-pyrazol-1-yl)benzenesulfonamide;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-82	 <p>4-(3,5-bis(4-methoxyphenyl)-1H-pyrazol-1-yl)benzenesulfonamide;</p>
B-83	 <p>4-(5-(4-chlorophenyl)-3-(4-methylphenyl)-1H-pyrazol-1-yl)benzenesulfonamide;</p>
B-84	 <p>4-(5-(4-chlorophenyl)-3-(4-nitrophenyl)-1H-pyrazol-1-yl)benzenesulfonamide;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-85	 <p>4-(5-(4-chlorophenyl)-3-(5-chloro-2-thienyl)-1H-pyrazol-1-yl)benzenesulfonamide;</p>
B-86	 <p>4-(4-chloro-3,5-diphenyl-1H-pyrazol-1-yl)benzenesulfonamide;</p>
B-87	 <p>4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-88	 <p>4-[5-phenyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;</p>
B-89	 <p>4-[5-(4-fluorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;</p>
B-90	 <p>4-[5-(4-methoxyphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-91	 <p>4-[5-(4-chlorophenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;</p>
B-92	 <p>4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;</p>
B-93	 <p>4-[4-chloro-5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide;</p>





<u>Compound Number</u>	<u>Structural Formula</u>
B-94	 <p data-bbox="573 688 1260 716">4-[3-(difluoromethyl)-5-(4-methylphenyl)-1H-pyrazol-1-yl]benzenesulfonamide;</p>
B-95	 <p data-bbox="594 1115 1260 1142">4-[3-(difluoromethyl)-5-phenyl-1H-pyrazol-1-yl]benzenesulfonamide;</p>
B-96	 <p data-bbox="573 1570 1260 1593">4-[3-(difluoromethyl)-5-(4-methoxyphenyl)-1H-pyrazol-1-yl]benzenesulfonamide;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-97	 <p>4-[3-cyano-5-(4-fluorophenyl)-1H-pyrazol-1-yl]benzenesulfonamide;</p>
B-98	 <p>4-[3-(difluoromethyl)-5-(3-fluoro-4-methoxyphenyl)-1H-pyrazol-1-yl] benzenesulfonamide;</p>
B-99	 <p>4-[5-(3-fluoro-4-methoxyphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl] benzenesulfonamide;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-100	 <p data-bbox="576 583 1156 617">4-[4-chloro-5-phenyl-1H-pyrazol-1-yl]benzenesulfonamide;</p>
B-101	 <p data-bbox="576 1003 1260 1031">4-[5-(4-chlorophenyl)-3-(hydroxymethyl)-1H-pyrazol-1-yl]benzenesulfonamide;</p>
B-102	 <p data-bbox="576 1545 1120 1604">4-[5-(4-(N,N-dimethylamino)phenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-103	 <p>5-(4-fluorophenyl)-6-[4-(methylsulfonyl)phenyl]spiro[2.4]hept-5-ene;</p>
B-104	 <p>4-[6-(4-fluorophenyl)spiro[2.4]hept-5-en-5-yl]benzenesulfonamide;</p>
B-105	 <p>6-(4-fluorophenyl)-7-[4-methylsulfonyl]phenyl]spiro[3.4]oct-6-ene;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-106	 <p>5-(3-chloro-4-methoxyphenyl)-6-[4-(methylsulfonyl)phenyl]spiro[2.4]hept-5-ene;</p>
B-107	 <p>4-[6-(3-chloro-4-methoxyphenyl)spiro[2.4]hept-5-en-5-yl]benzenesulfonamide;</p>
B-108	 <p>5-(3,5-dichloro-4-methoxyphenyl)-6-[4-(methylsulfonyl)phenyl]spiro[2.4]hept-5-ene;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-109	 <p>5-(3-chloro-4-fluorophenyl)-6-[4-(methylsulfonyl)phenyl]spiro[2.4]hept-5-ene;</p>
B-110	 <p>4-[6-(3,4-dichlorophenyl)spiro[2.4]hept-5-en-5-yl]benzenesulfonamide;</p>
B-111	 <p>2-(3-chloro-4-fluorophenyl)-4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)thiazole;</p>
B-112	 <p>2-(2-chlorophenyl)-4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)thiazole;</p>





<u>Compound Number</u>	<u>Structural Formula</u>
B-113	 <p>5-(4-fluorophenyl)-4-(4-methylsulfonylphenyl)-2-methylthiazole;</p>
B-114	 <p>4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-2-trifluoromethylthiazole;</p>
B-115	 <p>4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-2-(2-thienyl)thiazole;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-116	 <p>4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-2-benzylaminothiazole;</p>
B-117	 <p>4-(4-fluorophenyl)-5-(4-methylsulfonylphenyl)-2-(1-propylamino)thiazole;</p>
B-118	 <p>2-((3,5-dichlorophenoxy)methyl)-4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]thiazole;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-119	 <p>5-(4-fluorophenyl)-4-(4-methylsulfonylphenyl)-2-trifluoromethylthiazole;</p>
B-120	 <p>1-methylsulfonyl-4-[1,1-dimethyl-4-(4-fluorophenyl)cyclopenta-2,4-dien-3-yl]benzene;</p>
B-121	 <p>4-[4-(4-fluorophenyl)-1,1-dimethylcyclopenta-2,4-dien-3-yl]benzenesulfonamide;</p>



<u>Compound Number</u>	<u>Structural Formula</u>
B-122	 <p>5-(4-fluorophenyl)-6-[4-(methylsulfonyl)phenyl]spiro[2.4]hepta-4,6-diene;</p>
B-123	 <p>4-[6-(4-fluorophenyl)spiro[2.4]hepta-4,6-dien-5-yl]benzenesulfonamide;</p>
B-124	 <p>6-(4-fluorophenyl)-2-methoxy-5-[4-(methylsulfonyl)phenyl]-pyridine-3-carbonitrile;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-125	 <p>2-bromo-6-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]-pyridine-3-carbonitrile;</p>
B-126	 <p>6-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]-2-phenyl-pyridine-3-carbonitrile;</p>
B-127	 <p>4-[2-(4-methylpyridin-2-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl] benzenesulfonamide;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-128	 <p>4-[2-(5-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;</p>
B-129	 <p>4-[2-(2-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;</p>
B-130	 <p>3-[1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazol-2-yl]pyridine;</p>
B-131	 <p>2-[1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazol-2-yl]pyridine;</p>





<u>Compound Number</u>	<u>Structural Formula</u>
B-132	 <p>2-methyl-4-[1-[4-(methylsulfonyl)phenyl-4-(trifluoromethyl)]-1H-imidazol-2-yl]pyridine;</p>
B-133	 <p>2-methyl-6-[1-[4-(methylsulfonyl)phenyl-4-(trifluoromethyl)]-1H-imidazol-2-yl]pyridine;</p>
B-134	 <p>4-[2-(6-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;</p>





<u>Compound Number</u>	<u>Structural Formula</u>
B-135	 <p>2-(3,4-difluorophenyl)-1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazole;</p>
B-136	 <p>4-[2-(4-methylphenyl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;</p>
B-137	 <p>2-(4-chlorophenyl)-1-[4-(methylsulfonyl)phenyl]-4-methyl-1H-imidazole;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-138	 <p>2-(4-chlorophenyl)-1-[4-(methylsulfonyl)phenyl]-4-phenyl-1H-imidazole;</p>
B-139	 <p>2-(4-chlorophenyl)-4-(4-fluorophenyl)-1-[4-(methylsulfonyl)phenyl]-1H-imidazole;</p>



<u>Compound Number</u>	<u>Structural Formula</u>
B-140	 <p>2-(3-fluoro-4-methoxyphenyl)-1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazole;</p>
B-141	 <p>1-[4-(methylsulfonyl)phenyl]-2-phenyl-4-trifluoromethyl-1H-imidazole;</p>
B-142	 <p>2-(4-methylphenyl)-1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazole;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-143	 <p>4-[2-(3-chloro-4-methylphenyl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;</p>
B-144	 <p>2-(3-fluoro-5-methylphenyl)-1-[4-(methylsulfonyl)phenyl]-4-(trifluoromethyl)-1H-imidazole;</p>
B-145	 <p>4-[2-(3-fluoro-5-methylphenyl)-4-(trifluoromethyl)-1H-imidazole-1-yl]benzenesulfonamide;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-146	 <p>2-(3-methylphenyl)-1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazole;</p>
B-147	 <p>4-[2-(3-methylphenyl)-4-trifluoromethyl-1H-imidazol-1-yl]benzenesulfonamide;</p>
B-148	 <p>1-[4-(methylsulfonyl)phenyl]-2-(3-chlorophenyl)-4-trifluoromethyl-1H-imidazole</p>
B-149	 <p>4-[2-(3-chlorophenyl)-4-trifluoromethyl-1H-imidazol-1-yl]benzenesulfonamide;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-150	 <p>4-[2-phenyl-4-trifluoromethyl-1H-imidazol-1-yl]benzenesulfonamide;</p>
B-151	 <p>4-[2-(4-methoxy-3-chlorophenyl)-4-trifluoromethyl-1H-imidazol-1-yl]benzenesulfonamide;</p>
B-152	 <p>1-allyl-4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)-1H-pyrazole;</p>
B-153	 <p>4-[1-ethyl-4-(4-fluorophenyl)-5-(trifluoromethyl)-1H-pyrazol-3-yl]benzenesulfonamide;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-154	 <p data-bbox="576 808 1136 861">N-phenyl-4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)-1H-pyrazol-1-yl]acetamide;</p>
B-155	 <p data-bbox="576 1396 1079 1449">ethyl[4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)-1H-pyrazol-1-yl]acetate;</p>
B-156	 <p data-bbox="576 1827 1266 1858">4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-1-(2-phenylethyl)-1H-pyrazole;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-157	 <p>4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-1-(2-phenylethyl)-5-(trifluoromethyl)pyrazole;</p>
B-158	 <p>1-ethyl-4-(4-fluorophenyl)-3-[4-(methylsulfonyl)phenyl]-5-(trifluoromethyl)-1H-pyrazole;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-159	 <p>5-(4-fluorophenyl)-4-(4-methylsulfonylphenyl)-2-trifluoromethyl-1H-imidazole;</p>
B-160	 <p>4-[4-(methylsulfonyl)phenyl]-5-(2-thiophenyl)-2-(trifluoromethyl)-1H-imidazole;</p>
B-161	 <p>5-(4-fluorophenyl)-2-methoxy-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyridine;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-162	 <p>2-ethoxy-5-(4-fluorophenyl)-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyridine;</p>
B-163	 <p>5-(4-fluorophenyl)-4-[4-(methylsulfonyl)phenyl]-2-(2-propynyloxy)-6-(trifluoromethyl)pyridine;</p>
B-164	 <p>2-bromo-5-(4-fluorophenyl)-4-[4-(methylsulfonyl)phenyl]-6-(trifluoromethyl)pyridine;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-165	 <p>4-[2-(3-chloro-4-methoxyphenyl)-4,5-difluorophenyl]benzenesulfonamide;</p>
B-166	 <p>1-(4-fluorophenyl)-2-[4-methylsulfonyl]phenyl]benzene;</p>
B-167	 <p>5-difluoromethyl-4-(4-methylsulfonylphenyl)-3-phenylisoxazole;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-168	 <p>4-[3-ethyl-5-phenylisoxazol-4-yl]benzenesulfonamide;</p>
B-169	 <p>4-[5-difluoromethyl-3-phenylisoxazol-4-yl]benzenesulfonamide;</p>
B-170	 <p>4-[5-hydroxymethyl-3-phenylisoxazol-4-yl]benzenesulfonamide;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-171	 <p>4-[5-methyl-3-phenyl-isoxazol-4-yl]benzenesulfonamide;</p>
B-172	 <p>1-[2-(4-fluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;</p>
B-173	 <p>1-[2-(4-fluoro-2-methylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-174	 <p>1-[2-(4-chlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;</p>
B-175	 <p>1-[2-(2,4-dichlorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;</p>
B-176	 <p>1-[2-(4-trifluoromethylphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-177	 <p>1-[2-(4-methylthiophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;</p>
B-178	 <p>1-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;</p>
B-179	 <p>4-[2-(4-fluorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;</p>





<u>Compound Number</u>	<u>Structural Formula</u>
B-180	 <p>1-[2-(3-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]-4-(methylsulfonyl)benzene;</p>
B-181	 <p>4-[2-(4-chlorophenyl)-4,4-dimethylcyclopenten-1-yl]benzenesulfonamide;</p>
B-182	 <p>4-[2-(4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-183	 <p>4-[2-(4-chlorophenyl)cyclopenten-1-yl]benzenesulfonamide;</p>
B-184	 <p>1-[2-(4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;</p>
B-185	 <p>1-[2-(2,3-difluorophenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-186	 <p>4-[2-(3-fluoro-4-methoxyphenyl)cyclopenten-1-yl]benzenesulfonamide;</p>
B-187	 <p>1-[2-(3-chloro-4-methoxyphenyl)cyclopenten-1-yl]-4-(methylsulfonyl)benzene;</p>
B-188	 <p>4-[2-(3-chloro-4-fluorophenyl)cyclopenten-1-yl]benzenesulfonamide;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-189	 <p>4-[2-(2-methylpyridin-5-yl)cyclopenten-1-yl]benzenesulfonamide;</p>
B-190	 <p>ethyl 2-[4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]oxazol-2-yl]-2-benzylacetate;</p>
B-191	 <p>2-[4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]oxazol-2-yl]acetic acid;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-192	 <p>2-(tert-butyl)-4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]oxazole;</p>
B-193	 <p>4-(4-fluorophenyl)-5-[4-(methylsulfonyl)phenyl]-2-phenyloxazole;</p>
B-194	 <p>4-(4-fluorophenyl)-2-methyl-5-[4-(methylsulfonyl)phenyl]oxazole;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-195	 <p>4-[5-(3-fluoro-4-methoxyphenyl)-2-trifluoromethyl-4-oxazolyl]benzenesulfonamide;</p>
B-196	 <p>6-chloro-7-(1,1-dimethylethyl)-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-197	 <p>6-chloro-8-methyl-2-trifluoromethyl-2H-1-benzopyran-3-carboxylic acid;</p>
B-198	 <p>5,5-dimethyl-3-(3-fluorophenyl)-4-methylsulfonyl-2(5H)-furanone;</p>





<u>Compound Number</u>	<u>Structural Formula</u>
B-199	 <p>6-chloro-2-trifluoromethyl-2H-1-benzothiopyran-3-carboxylic acid;</p>
B-200	 <p>4-[5-(4-chlorophenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;</p>
B-201	 <p>4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;</p>


<u>Compound Number</u>	<u>Structural Formula</u>
B-202	 <p>4-[5-(3-fluoro-4-methoxyphenyl)-3-(difluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide;</p>
B-203	 <p>3-[1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazol-2-yl]pyridine;</p>
B-204	 <p>2-methyl-5-[1-[4-(methylsulfonyl)phenyl]-4-trifluoromethyl-1H-imidazol-2-yl]pyridine;</p>




<u>Compound Number</u>	<u>Structural Formula</u>
B-205	 <p>4-[2-(5-methylpyridin-3-yl)-4-(trifluoromethyl)-1H-imidazol-1-yl]benzenesulfonamide;</p>
B-206	 <p>4-[5-methyl-3-phenylisoxazol-4-yl]benzenesulfonamide;</p>
B-207	 <p>4-[5-hydroxymethyl-3-phenylisoxazol-4-yl]benzenesulfonamide;</p>





<u>Compound Number</u>	<u>Structural Formula</u>
B-208	 <p data-bbox="560 777 1250 808">[2-trifluoromethyl-5-(3,4-difluorophenyl)-4-oxazolyl]benzenesulfonamide;</p>
B-209	 <p data-bbox="560 1260 1136 1302">4-[2-methyl-4-phenyl-5-oxazolyl]benzenesulfonamide;</p>
B-210	 <p data-bbox="560 1774 1250 1816">4-[5-(2-fluoro-4-methoxyphenyl)-2-trifluoromethyl-4-oxazolyl]benzenesulfonamide;</p>






<u>Compound Number</u>	<u>Structural Formula</u>
B-211	 <chem>CC1=CC=C(C(=C1)C(=O)O)Nc2cc(Cl)ccc2S(=O)(=O)Nc3cc(F)ccc3</chem>
B-212	 <i>N</i> -(4-nitro-2-phenoxy-phenyl)-methanesulfonamide <chem>CO=S(=O)(=O)Nc1cc(OC2=CC=CC=C2)ccc1[N+](=O)[O-]</chem>
B-213	 <i>N</i> -[6-(2,4-difluoro-phenoxy)-1-oxo-inden-5-yl]-methanesulfonamide <chem>CO=S(=O)(=O)Nc1ccc2c(c1)c(c3ccccc23)C(=O)Oc4cc(F)ccc4F</chem>






<u>Compound Number</u>	<u>Structural Formula</u>
B-214	 <p data-bbox="570 737 1258 783"><i>N</i>-[6-(2,4-difluoro-phenylsulfanyl)-1-oxo-1<i>H</i>-inden-5-yl]-methanesulfonamide, sodium salt</p>
B-215	 <p data-bbox="570 947 1133 972"><i>N</i>-[5-(4-fluoro-phenylsulfanyl)-thiophen-2-yl]-methanesulfonamide</p>
B-216	 <p data-bbox="570 1419 1219 1476">3-(3,4-difluoro-phenoxy)-4-(4-methanesulfonyl-phenyl)-5-methyl-5-(2,2,2-trifluoro-ethyl)-5<i>H</i>-furan-2-one</p>
B-217	 <p data-bbox="570 1860 1258 1911">(5<i>Z</i>)-2-amino-5-[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene]-4(5<i>H</i>)-thiazolone</p>






<u>Compound Number</u>	<u>Structural Formula</u>
B-218	CS-502
B-219	LAS-34475
B-220	LAS-34555
B-221	S-33516
B-222	SD-8381
B-223	L-783003
B-224	 <p>N-[3-(formylamino)-4-oxo-6-phenoxy-4H-1-benzopyran-7-yl]-methanesulfonamide</p>
B-225	D-1367
B-226	L-748731


<u>Compound Number</u>	<u>Structural Formula</u>
B-227	 <p>(6aR,10aR)-3-(1,1-dimethylheptyl)-6a,7,10,10a-tetrahydro-1-hydroxy-6,6-dimethyl-1-6H-dibenzo[b,d]pyran-9-carboxylic acid</p>
B-228	CGP-28238
B-229	 <p>4-[[3,5-bis(1,1-dimethylethyl)-4-hydroxyphenyl]methylene]dihydro-2-methyl-2H-1,2-oxazin-3(4H)-one</p>
B-230	GR-253035
B-231	 <p>2-(6-dioxo-9H-purin-8-yl)cinnamic acid</p>
B-232	S-2474

<u>Compound Number</u>	<u>Structural Formula</u>
B-233	
B-234	
B-235	
B-236	

<u>Compound Number</u>	<u>Structural Formula</u>
B-237	 <chem>COc1ccc(cc1)C2=C(C(=O)N2c3cc(F)cc(F)c3)c4cc(O)cc(O)c4</chem>
B-238	 <chem>COc1ccc(cc1)C2=C(C(=O)N2c3cc(Cl)cc(O)c3)c4cc(O)cc(O)c4</chem>
B-239	 <chem>COc1ccc(cc1)C2=C(C(=O)N2c3cc(F)cc(O)c3)c4cc(O)cc(O)c4</chem>
B-240	 <chem>COc1ccc(cc1)C2=C(C(=O)N2c3cc(C)cc(O)c3)c4cc(O)cc(O)c4</chem>
B-241	 <chem>COc1ccc(cc1)C2=C(C(=O)N2c3cc(C(F)(F)F)cc(O)c3)c4cc(O)cc(O)c4</chem>

<u>Compound Number</u>	<u>Structural Formula</u>
B-242	
B-243	
B-244	
B-245	
B-246	

<u>Compound Number</u>	<u>Structural Formula</u>
B-247	 <chem>COc1ccc(cc1)C2=C(C(=O)N2c3cc(F)cc(F)c3)C4=CC=C(CS(=O)(=O)C4)C5=CC=CC=C5</chem>
B-248	 <chem>CC1=CC=C(N1C2=CC(=O)C(C2)c3ccc(F)cc3)C4=CC=C(CS(=O)(=O)C4)C5=CC=CC=C5</chem>
B-249	 <chem>COc1cccc(N1C2=CC(=O)C(C2)c3ccc(F)cc3)C4=CC=C(CS(=O)(=O)C4)C5=CC=CC=C5</chem>
B-250	 <chem>CC1=CC=C(N1C2=CC(=O)C(C2)c3ccc(F)cc3)C4=CC=C(CS(=O)(=O)C4)C5=CC=C(C5)C(F)(F)F</chem>
B-251	 <chem>COc1ccc(cc1)C2=C(C(=O)N2c3cc(F)ccc3)C4=CC=C(CS(=O)(=O)C4)C5=CC=CC=C5</chem>

<u>Compound Number</u>	<u>Structural Formula</u>
B-252	

[0415] The cyclooxygenase-2 selective inhibitor employed in the present invention can exist in tautomeric, geometric or stereoisomeric forms. Generally speaking, suitable cyclooxygenase-2 selective inhibitors that are in tautomeric, geometric or stereoisomeric forms are those compounds that inhibit cyclooxygenase-2 activity by about 25%, more typically by about 50%, and even more typically, by about 75% or more when present at a concentration of 100 μ M or less. The present invention contemplates all such compounds, including cis- and trans-geometric isomers, E- and Z-geometric isomers, R- and S-enantiomers, diastereomers, d-isomers, l-isomers, the racemic mixtures thereof and other mixtures thereof. Pharmaceutically acceptable salts of such tautomeric, geometric or stereoisomeric forms are also included within the invention. The terms "cis" and "trans", as used herein, denote a form of geometric isomerism in which two carbon atoms connected by a double bond will each have a hydrogen atom on the same side of the double bond ("cis") or on opposite sides of the double bond ("trans"). Some of the compounds described contain alkenyl groups, and are meant to include both cis and trans or "E" and "Z" geometric forms. Furthermore, some of the compounds described contain one or more stereocenters and are meant to include R, S, and mixtures or R and S forms for each stereocenter present.

[0416] The cyclooxygenase-2 selective inhibitors utilized in the present invention may be in the form of free bases or pharmaceutically acceptable acid addition salts thereof. The term "pharmaceutically-acceptable salts" are salts commonly used to form alkali metal salts and to form addition salts of free acids or free bases. The nature of the salt may vary, provided that it is pharmaceutically acceptable. Suitable pharmaceutically acceptable acid addition salts of compounds for use in the present methods may be prepared from an inorganic acid or from an organic acid. Examples of such inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, carbonic, sulfuric and phosphoric acid. Appropriate organic acids may be selected from aliphatic, cycloaliphatic, aromatic,

araliphatic, heterocyclic, carboxylic and sulfonic classes of organic acids, examples of which are formic, acetic, propionic, succinic, glycolic, gluconic, lactic, malic, tartaric, citric, ascorbic, glucuronic, maleic, fumaric, pyruvic, aspartic, glutamic, benzoic, anthranilic, mesylic, 4-hydroxybenzoic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, benzenesulfonic, pantothenic, 2-hydroxyethanesulfonic, toluenesulfonic, sulfanilic, cyclohexylaminosulfonic, stearic, algenic, hydroxybutyric, salicylic, galactaric and galacturonic acid. Suitable pharmaceutically-acceptable base addition salts of compounds of use in the present methods include metallic salts made from aluminum, calcium, lithium, magnesium, potassium, sodium and zinc or organic salts made from N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine) and procaine. All of these salts may be prepared by conventional means from the corresponding compound by reacting, for example, the appropriate acid or base with the compound of any Formula set forth herein.

[0417] The cyclooxygenase-2 selective inhibitors of the present invention can be formulated into pharmaceutical compositions and administered by a number of different means that will deliver a therapeutically effective dose. Such compositions can be administered orally, parenterally, by inhalation spray, rectally, intradermally, transdermally, or topically in dosage unit formulations containing conventional nontoxic pharmaceutically acceptable carriers, adjuvants, and vehicles as desired. Topical administration may also involve the use of transdermal administration such as transdermal patches or iontophoresis devices. The term parenteral as used herein includes subcutaneous, intravenous, intramuscular, or intrasternal injection, or infusion techniques. Formulation of drugs is discussed in, for example, Hoover, John E., *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, Pennsylvania (1975), and Liberman, H.A. and Lachman, L., Eds., *Pharmaceutical Dosage Forms*, Marcel Dekker, New York, N.Y. (1980).

[0418] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions, can be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed, including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are useful in the preparation of injectables. Dimethyl acetamide, surfactants including ionic and non-ionic detergents, and

polyethylene glycols can be used. Mixtures of solvents and wetting agents such as those discussed above are also useful.

[0419] Suppositories for rectal administration of the compounds discussed herein can be prepared by mixing the active agent with a suitable non-irritating excipient such as cocoa butter, synthetic mono-, di-, or triglycerides, fatty acids, or polyethylene glycols which are solid at ordinary temperatures but liquid at the rectal temperature, and which will therefore melt in the rectum and release the drug.

[0420] Solid dosage forms for oral administration may include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the compounds are ordinarily combined with one or more adjuvants appropriate to the indicated route of administration. If administered *per os*, the compounds can be admixed with lactose, sucrose, starch powder, cellulose esters of alkanolic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets can contain a controlled-release formulation as can be provided in a dispersion of active compound in hydroxypropylmethyl cellulose. In the case of capsules, tablets, and pills, the dosage forms can also comprise buffering agents such as sodium citrate, or magnesium or calcium carbonate or bicarbonate. Tablets and pills can additionally be prepared with enteric coatings.

[0421] For therapeutic purposes, formulations for parenteral administration can be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions can be prepared from sterile powders or granules having one or more of the carriers or diluents mentioned for use in the formulations for oral administration. The compounds can be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art.

[0422] Liquid dosage forms for oral administration can include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, and elixirs containing inert diluents commonly used in the art, such as water. Such compositions can also comprise adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring, and perfuming agents.

[0423] The amount of active ingredient that can be combined with the carrier materials to produce a single dosage of the cyclooxygenase-2 selective inhibitor will vary

depending upon the patient and the particular mode of administration. In general, the pharmaceutical compositions may contain a cyclooxygenase-2 selective inhibitor in the range of about 0.1 to 2000 mg, more typically, in the range of about 0.5 to 500 mg and still more typically, between about 1 and 200 mg. A daily dose of about 0.01 to 100 mg/kg body weight, or more typically, between about 0.1 and about 50 mg/kg body weight and even more typically, from about 1 to 20 mg/kg body weight, may be appropriate. The daily dose is generally administered in one to about four doses per day.

[0424] In one embodiment, when the cyclooxygenase-2 selective inhibitor comprises rofecoxib, it is typical that the amount used is within a range of from about 0.15 to about 1.0 mg/day·kg, and even more typically, from about 0.18 to about 0.4 mg/day·kg.

[0425] In still another embodiment, when the cyclooxygenase-2 selective inhibitor comprises etoricoxib, it is typical that the amount used is within a range of from about 0.5 to about 5 mg/day·kg, and even more typically, from about 0.8 to about 4 mg/day·kg.

[0426] Further, when the cyclooxygenase-2 selective inhibitor comprises celecoxib, it is typical that the amount used is within a range of from about 1 to about 20 mg/day·kg, even more typically, from about 1.4 to about 8.6 mg/day·kg, and yet more typically, from about 2 to about 3 mg/day·kg.

[0427] When the cyclooxygenase-2 selective inhibitor comprises valdecoxib, it is typical that the amount used is within a range of from about 0.1 to about 5 mg/day·kg, and even more typically, from about 0.8 to about 4 mg/day·kg.

[0428] In a further embodiment, when the cyclooxygenase-2 selective inhibitor comprises parecoxib, it is typical that the amount used is within a range of from about 0.1 to about 5 mg/day·kg, and even more typically, from about 1 to about 3 mg/day·kg.

[0429] Those skilled in the art will appreciate that dosages may also be determined with guidance from Goodman & Goldman's The Pharmacological Basis of Therapeutics, Ninth Edition (1996), Appendix II, pp. 1707-1711 and from Goodman & Goldman's The Pharmacological Basis of Therapeutics, Tenth Edition (2001), Appendix II, pp. 475-493.

SEROTONIN-MODULATING AGENTS

[0430] In addition to a cyclooxygenase-2 selective inhibitor, the composition of the invention also comprises a therapeutically effective amount of a serotonin-modulating agent or a pharmaceutically acceptable salt or prodrug thereof. A number of serotonin-modulating agents may be employed in the present invention. In some aspects, the

serotonin-modulating agent may reverse or lessen central nervous system cell damage following a reduction in blood flow to the central nervous system. In other aspects, the serotonin-modulating agent may reverse or lessen central nervous system cell damage following a traumatic brain or spinal cord injury.

[0431] In one aspect of the invention, the serotonin-modulating agent is a serotonin receptor antagonist. In one embodiment, the serotonin receptor antagonist is a 5-HT₁ antagonist. In one alternative of this embodiment, the 5-HT₁ antagonist is selected from the group consisting of:

- [0432] 3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol;
- [0433] 4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile;
- [0434] 3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide;
- [0435] N-[4-methoxy-3-(4-methyl-1-piperazinyl)phenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-1,1'-biphenyl-4-carboxamide hydrochloride;
- [0436] 1-[(1-methylethyl)amino]-3-[2-(1H-pyrrol-1-yl)phenoxy]propan-2-ol;
- [0437] 1-(2-methoxyphenyl)-4-(4-succinimidobutyl)piperazine;
- [0438] 1-(2-methoxyphenyl)-4-(4-phthalimidobutyl)piperazine;
- [0439] 1-(1H-indol-4-yloxy)-3-[(1-methylethyl)amino]-2-propanol;
- [0440] (S)-1-(1H-indol-4-yloxy)-3-[(1-methylethyl)amino]-2-propanol;
- [0441] N-[3-[3-(dimethylamino)ethoxy]-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-[1,1'-biphenyl]-4-carboxamide;
- [0442] 8-[(2,3-dihydro-1,4-benzodioxin-2-yl)methyl]-1-phenyl-1,3,8-triazaspiro[4,5]decan-4-one; and
- [0443] (S)-N-tert-butyl-3-(4-(2-methoxyphenyl)-piperazin-1-yl)-2-phenylpropanamide;
- [0444] or a pharmaceutically acceptable salt or prodrug thereof.
- [0445] In another embodiment, the serotonin receptor antagonist is a 5-HT₂ antagonist. In one alternative of this embodiment, the 5-HT₂ antagonist is selected from the group consisting of:
- [0446] 8-[3-(4-fluorophenoxy)propyl]-1-phenyl-1,3,8-triazaspiro[4.5]-decan-4-one;
- [0447] N-[2-[[3-(dimethylamino)propyl]thio]phenyl]-3-phenyl-2-propenamide;
- [0448] 4-(5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-methylpiperidine;
- [0449] 8-chloro-11-(1-piperazinyl)-5H-dibenzo[b,e][1,4]diazepine;

- [0450] 4-(4-fluorobenzoyl)-1-(4-phenylbutyl)-piperidine oxalate;
- [0451] 3-[2-[4-(4-fluorobenzoyl)-1-piperidinyl]ethyl]-2,4[1H,3H]-quinazolinedione;
- [0452] α -phenyl-1-(2-phenylethyl)-4-piperidinemethanol;
- [0453] metergoline phenylmethyl ester;
- [0454] 1,2,3,4,10,14b-hexahydro-2-methyldibenzo[c,f]pyrazino[1,2-a]azepine;
- [0455] 8-[5-(2,4-dimethoxy-5-(4-trifluoromethylphenylsulphonamido) phenyl-5-oxopentyl]-1,3,8-triazaspiro[4.5]decane-2,4-dione;
- [0456] N-(1-methyl-1H-indol-5-yl)-N'-3-pyridinyl-urea;
- [0457] N-(1-methyl-5-indolyl)-N'-(3-methyl-5-isothiazolyl)urea;
- [0458] (+)-cis-4,5,7a,8,9,10,11,11a-octahydro-7H-10-methylindolo[1,7-bc][2,6]-naphthyridine; and
- [0459] 8-[4-(4-fluorophenyl)-4-oxobutyl]-1-phenyl-1,3,8-triazaspiro[4,5]decan-4-one;
- [0460] or a pharmaceutically acceptable salt or prodrug thereof.
- [0461] In yet another embodiment, the serotonin receptor antagonist is a 5-HT₃ antagonist. In one alternative of this embodiment, the 5-HT₃ antagonist is selected from the group consisting of:
- [0462] 3aS-2-[(S)-1-azabicyclo [2.2.2] oct-3-yl]-2,3,3a,4,5,6-hexahydro-1-oxo-1H-benz [de]isoquinoline hydrochloride;
- [0463] 3-(4-allylpiperazin-1-yl)-2-quinoxalinecarbonitrile maleate;
- [0464] proparyl 3,5-dichlorobenzoate;
- [0465] proparyl 3,5-dimethylbenzoate; and
- [0466] N-(1-azabicyclo[2.2.2]oct-3-yl)-6-chloro-4-methyl-3-oxo-3,4-dihydro-2H-1,4-benzoxazine-8-carboxamide;
- [0467] or a pharmaceutically acceptable salt or prodrug thereof.
- [0468] In a further embodiment, the serotonin receptor antagonist is a 5-HT₄ antagonist. In one alternative of this embodiment, the 5-HT₄ antagonist is selected from the group consisting of:
- [0469] 3-methyl-1H-indole-3-carboxylic acid;
- [0470] 3-(piperidin-1-yl)propyl-4-amino-5-chloro-2-methoxybenzoate;
- [0471] 1-[4-amino-5-chloro-2-(3,5-dimethoxyphenyl)methoxy]-3-[1-[2-methylsulphonylamino]ethyl]piperidin-4-yl]propan-1-one; and
- [0472] 1-piperidinylethyl-1H-indole-3-carboxylate;
- [0473] or a pharmaceutically acceptable salt or prodrug thereof.

[0474] In another embodiment, the serotonin receptor antagonist is a 5-HT₆ antagonist. In one alternative of this embodiment, the 5-HT₆ antagonist is metergoline phenylmethyl ester, or a pharmaceutically acceptable salt or prodrug thereof.

[0475] In yet another embodiment, the serotonin receptor antagonist is a 5-HT₇ antagonist. In one alternative of this embodiment, the 5-HT₇ antagonist is selected from the group consisting of:

[0476] metergoline phenylmethyl ester, and

[0477] 1-[1-[4,4-bis(4-fluorophenyl)butyl]-4-piperidiny]1,3-dihydro-2H-benzimidazol-2-one, or a pharmaceutically acceptable salt or prodrug thereof.

[0478] In another aspect of the invention, the serotonin-modulating agent is a serotonin receptor agonist. In one embodiment, the serotonin receptor agonist is a 5-HT₁ agonist. In one alternative of this embodiment, the 5-HT₁ agonist is selected from the group consisting of:

[0479] 6-chloro-2-[piperidiny-4-thio]pyridine;

[0480] 8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione;

[0481] 1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine;

[0482] 5-hydroxy-3-(1-methylpiperidin-4-yl)-1H-indole;

[0483] 8-[4-[4-(2-pyrimidinyl)-1-piperazinyl]butyl]-8-azaspiro[4.5]decane-7,8-dione;

[0484] 5-carboxamidotryptamine maleate;

[0485] 7-trifluoromethyl-4-(4-methyl-1-piperazinyl)pyrrolo[1,2-a]-quinoxaline;

[0486] 1,4-dihydro-3-(1,2,3,6-tetrahydro-4-pyridinyl)-5H-pyrrolo[3,2-b]pyridin-5-one;

[0487] 5-propoxy-3-(1,2,3,6-tetrahydro-4-pyridinyl)-1H-pyrrolo[3,2-b]pyridine hydrochloride;

[0488] 3-[3-(2-dimethylaminoethyl)-1H-indol-5-yl]-N-(4-methoxybenzyl)acrylamide;

[0489] 8-hydroxy-2-dipropylaminotetralin hydrobromide;

[0490] (2R)-(+)-8-hydroxy-2-(di-n-propylamino)tetralin;

[0491] (RS)-trans-8-hydroxy-2-[N-n-propyl-N-(3'-iodo-2'-propenyl)amino]tetralin;

[0492] 2-[5-[3-(-4-methylsulphonylamino)benzyl]-1,2,4-oxadiazol-5-yl]-1H-indol-3-yl]ethanamine;

[0493] 8-[2-(1,4-benzodioxan-2-ylmethylamino)ethyl]-8-azaspiro[4.5]decane-7,9-dione;

[0494] nonyloxytryptamineoxalate;

[0495] 5-methoxy-3-(1,2,5,6-tetrahydro-4-pyridinyl)-1H-indole; and

[0496] N-(3-trifluoromethylphenyl)piperazine;

[0497] or a pharmaceutically acceptable salt or prodrug thereof.

[0498] In another embodiment, the serotonin receptor agonist is a 5-HT₂ agonist.

In one alternative of this embodiment, the 5-HT₂ agonist is selected from the group consisting of:

[0499] α -methyl-5-(2-thienylmethoxy)-1H-indole-3-ethanamine;

[0500] 1-(3-chlorophenyl)piperazine;

[0501] α -methyl-5-hydroxytryptamine maleate; and

[0502] 6-chloro-2-(1-piperazinyl)pyrazine;

[0503] or a pharmaceutically acceptable salt or prodrug thereof.

[0504] In yet another embodiment, the serotonin receptor agonist is a 5-HT₃ agonist. In one alternative of this embodiment, the 5-HT₃ agonist is selected from the group consisting of:

[0505] 1-(3-chlorophenyl)biguanide;

[0506] 2-methyl-5-hydroxytryptamine hydrochloride;

[0507] 2-(1-N-methylpiperazinyl)quinoline;

[0508] 1-phenylbiguanide hydrochloride;

[0509] 2-(1-piperazinyl)quinoline;

[0510] (R)-N-(1-azabicyclo[2.2.2]oct-3-yl)-2-(1-methyl-1H-indol-3-yl)-2-(1-methyl-1H-indol-3-yl)-2-oxoacetamide; and

[0511] 1-(6-chloro-2-pyridinyl)-4-piperidinamine;

[0512] or a pharmaceutically acceptable salt or prodrug thereof.

[0513] In a further embodiment, the serotonin receptor agonist is a 5-HT₄ agonist. In one alternative of this embodiment, the 5-HT₄ agonist is selected from the group consisting of:

[0514] 2-[1-(4-piperonyl)piperazinyl]benzothiazole;

[0515] 1-(4-amino-5-chloro-2-methoxyphenyl)-3-[1-butyl-4-piperidinyl]-1-propanone; and

[0516] 1-(4-amino-5-chloro-2-methoxyphenyl)-3-[1-2-methylsulphonylamino)ethyl-4-piperidinyl]-1-propanone;

[0517] or a pharmaceutically acceptable salt or prodrug thereof.

[0518] In a still further embodiment, the serotonin receptor agonist is a 5-HT₅ agonist. In one alternative of this embodiment, the 5-HT₅ agonist is 5-carboxamidotryptamine maleate, or a pharmaceutically acceptable salt or prodrug thereof.

[0519] In another embodiment, the serotonin receptor agonist is a 5-HT₆ agonist. In one alternative of this embodiment, the 5-HT₆ agonist is 2-methyl-5-hydroxytryptamine hydrochloride, or a pharmaceutically acceptable salt or prodrug thereof.

[0520] In yet another embodiment, the serotonin receptor agonist is a 5-HT₇ agonist. In one alternative of this embodiment, the 5-HT₇ agonist is selected from the group consisting of 5-carboxamidotryptamine maleate, and (+,-)-8-hydroxy-2-dipropylaminotetralin, or a pharmaceutically acceptable salt or prodrug thereof.

[0521] In still another embodiment, compounds that are useful for the serotonin-modulating agent or a pharmaceutically acceptable salt or prodrug thereof in connection with the present invention, include, but are not limited to:

[0522] N-acetyltryptamine;

[0523] p-chlorophenylalanine;

[0524] 8-chloro-11-(4-methyl-1-piperazinyl)-5H-dibenzo[b,e][1,4]diazepine;

[0525] (5' α ,10 α)-9,10-dihydro-12'-hydroxy-2'-(1-methylethyl)-5'-(phenylmethyl)-ergotaman-3',6',18-trione;

[0526] 9,10-dihydro-12'-hydroxy-2'-methyl-5'-(phenylmethyl)ergotaman-3',6',18-trione;

[0527] 1-[10,11-dihydro-8-(methylthio)dibenzo[b,f]thiepin-10-yl]-4-methylpiperazine;

[0528] [8 β (S)]-9,10-didehydro-N-[1-(hydroxymethyl)propyl]-6-methylergoline-8-carboxamide;

[0529] [8 β (S)]-9,10-didehydro-N-[1-(hydroxymethyl)propyl]-1,6-dimethylergoline-8-carboxamide;


[0530] cis-9-octadecenoamide;

[0531] [1aR-(1aR*,4E,7aS*,10aS*,-10bR*)]-2,3-6,7,7a,8,10a,10b-octahydro-1a,5-dimethyl-8-methyleneoxireno[9,10]cyclodeca[1,2-b]furan-9-(1aH)-one; and

[0532] N-(3-trifluoromethylphenyl)piperazine.

[0533] In a further embodiment, the serotonin-modulating agent is selected from the group consisting of compounds having the general Formula I shown below and containing, by way of example and not limitation, the compounds listed below. Furthermore, the serotonin-modulating agents useful in the practice of the present

invention are described in U.S. Patent No. 5,436,246 which is herein incorporated by reference in its entirety.



Formula I

[0534] wherein:

[0535] Y is hydrogen or C₁₋₃ alkyl;

[0536] R is hydrogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, halogen, -CF₃, -OCF₃, or -OH;

[0537] R₁ is hydrogen, cycloalkyl, C₁₋₆ alkyl, phenyl optionally substituted, phenylalkyl, or phenylamidoalkyl;

[0538] X is hydrogen, -(CH₂)_nX₁, -CH=CHX₁ or -CHX₂-(CH₂)_q-CH₃;

[0539] wherein:

[0540] n is an integer from 0-2;

[0541] q is the integer 0 or 1;

[0542] X₁ is -OH, -OR₂, -NR₂R₃, -CO₂R₂, -CONR₂R₃, -CN, CH₂OH, or -COR₂;

[0543] R₂ and R₃ are each independently hydrogen, C₁₋₄ alkyl, phenyl optionally substituted, phenylalkyl, or R₂ and R₃ together form a (CH₂)_m cycloalkyl, where m is an integer from 2-6; and X₂ is -OR₄ or -NR₄R₅ in which R₄ and R₅ are each independently hydrogen or C₁₋₄ alkyl, provided however, when n is 0 or X is -CH=CHX₁, then X₁ is not OH, OR₂, or NR₂R₃.

[0544] Examples of suitable compounds having formula I include:

[0545] 4-[4-(2-phenylethyl)-1-piperazinyl]-benzo[b]thiophene-2-methanolmonohydrochloride;

[0546] 4-[4-(2-phenylethyl)-1-piperazinyl]-benzo[b]thiophene-2-carboxamide;

[0547] 4-[4-(2-phenylethyl)-1-piperazinyl]-benzo[b]thiophene-2-nitrile;

- [0548] 4-[4-(3-phenylpropyl)-1-piperazinyl]-benzo[b]thiophene-2-methanol;
- [0549] 4-[4-(3-phenylpropyl)-1-piperazinyl]-benzo[b]thiophene-2-carboxamide;
- [0550] 4-[4-[2-(4-methoxyphenyl)ethyl]-1-piperazinyl]-benzo[b]thiophene-2-methanol;
- [0551] 4-[4-[2-(4-chlorophenyl)ethyl]-1-piperazinyl]-benzo[b]thiophene-2-carboxamide;
- [0552] 4-[4-[2-(4-chlorophenyl)ethyl]-1-piperazinyl]-benzo[b]thiophene-2-methanol;
- [0553] 4-[4-[2-(4-methylphenyl)ethyl]-1-piperazinyl]-benzo[b]thiophene-2-methanol;
- [0554] 4-[4-(2-phenylethyl)-1-piperazinyl]-benzo[b]thiophene-2-(N-methyl)-carboxamide;
- [0555] 4-[4-(2-phenylethyl)-1-piperazinyl]-benzo[b]thiophene-2-(N,N-dimethyl)-carboxamide;
- [0556] 4-[4-[2-(4-methylphenyl)ethyl]-1-piperazinyl]-benzo[b]thiophene-2-carboxamide;
- [0557] 4-[4-[2-(4-fluorophenyl)ethyl]-1-piperazinyl]-benzo[b]thiophene-2-methanol;
- [0558] 4-[4-[2-(4-fluorophenyl)ethyl]-1-piperazinyl]-benzo[b]thiophene-2-carboxamide;
- [0559] ethyl-4-[(4-propyl)-1-piperazinyl]benzo[b]thiophene-2-carboxylatehydrochloride;
- [0560] 4-[(4-propyl)-1-piperazinyl]benzo[b]thiophene-2-methanolhydrochloride;
- [0561] 4-[4-(2-phenylethyl)-1-piperazinyl]-benzo[b]thiophene-2-(N-ethyl)carboxamidehydrochloride;
- [0562] 4-[4-(2-phenylethyl)-1-piperazinyl]-benzo[b]thiophene-2-(O-methyl)-methanolhydrochloride;
- [0563] 4-[4-propyl-1-piperazinyl]-benzo[b]thiophene-2-[N-methyl]carboxamidehydrochloride;
- [0564] 4-[4-methyl-1-piperazinyl]-benzo[b]thiophene-2-methanolhydrochloride;
- [0565] 4-[4-(2-phenylethyl)-1-piperazinyl]-benzo[b]thiophene-2-(N-methyl-N-methoxy)-carboxamidehydrochloride;
- [0566] 2-[4-[4-(2-phenylethyl)-1-piperazinyl]benzo[b]thiophene-2-](2-propanol)hydrochloride hemihydrate;

- [0567] 1-[4-(4-phenethyl-piperazin-1-yl)-benzo[b]thiophen-2-yl]-ethanonehydrochloride;
- [0568] 1-[4-(4-phenethyl-piperazin-1-yl)-benzo[b]thiophen-2-yl]-ethanolhydrochloride;
- [0569] 4-[4-phenylmethyl-1-piperazinyl]-benzo[b]thiophene-2-methoxymethylhydrochloride;
- [0570] 4-(1-piperazinyl)-benzo[b]thiophene-2-methoxymethylhydrochloride;
- [0571] 4-[4-(2-(4-fluorophenyl)-ethyl)-1-piperazinyl]benzo[b]thiophene-2-methoxymethylhydrochloride;
- [0572] 4-[4-(2-phenylethyl)-1-piperazinyl]-benzo[b]thiophene-2-carboxaldehyde;
- [0573] 4-[4-(4-phenylcarbomoyl-butyl)-piperazin-1-yl]-benzo[b]thiophene-2-carboxylicacidethylesterhydrochloride;
- [0574] 4-(1-piperazinyl)benzo[b]thiophene-2-(N-methyl)carboxamide;
- [0575] 4-4-[2-(4-nitrophenyl)ethyl]-1-piperazinyl-benzo[b]thiophene-2-methanolhydrochloridedihydrochloride;
- [0576] 4-(1-piperazinyl)benzo[b]thiophene-2-methanolhydrochloride;
- [0577] ethyl-4-[4-[2-(4-nitrophenyl)ethyl]-1-piperazinyl-benzo[b]thiophene-2-carboxylatehydrochloride;
- [0578] 5-[4-(2-Hydroxymethyl-benzo[b]thiophen-4-yl)-piperazin-1-yl]-pentanoicacidphenylamidehydrochloride;
- [0579] 2-[4-(4-phenethyl-piperazin-1-yl)-benzo[b]thiophen-2-ylmethyl]-isoindole-1,3-dionehydrochloride;
- [0580] 4-[4-(2-phenylethyl)-1-piperazinyl]-benzo[b]thiophene-2-methanamedihydrochloride;
- [0581] [4-(4-phenethyl-piperazin-1-yl)-benzo[b]thiophen-2-yl]-piperidin-1-ylmethanonehydrochloride;
- [0582] [4-(4-phenethyl-piperazin-1-yl)-benzo[b]thiophen-2-yl]pyrrolidin-1-ylmethanonehydrochloride;
- [0583] 3-[4-(4-phenethyl-piperazin-1-yl)-benzo[b]thiophen-2-yl]-acrylicacidethylesterhydrochloride-1-yl]-acrylicacidethylesterhydrochloride;
- [0584] 3-[4-(4-phenethyl-piperazin-1-yl)-benzo[b]thiophen-2-yl]-prop-2-en-1-olhydrochloride;
- [0585] 3-[4-(4-phenethyl-piperazin-1-yl)-benzo[b]thiophen-2-yl]-acrylonitrilehydrochloride;

[0586] 3-[4-(4-phenethyl-piperazin-1-yl)-benzo[b]thiophen-2-yl]-acrylamidehydrochloride;


[0587] 3-[4-(4-phenethyl-piperazin-1-yl)-benzo[b]thiophen-2-yl]-propionicacidethylesterhydrochloride;

[0588] 3-[4-(4-phenethyl-piperazin-1-yl)-benzo[b]thiophen-2-yl]-propan-1-olhydrochloride;

[0589] 3-[4-(4-phenethyl-piperazin-1-yl)-benzo[b]thiophen-2-yl]-propionitrilehydrochloride; and

[0590] 3-[4-(4-phenethyl-piperazin-1-yl)-benzo[b]thiophen-2-yl]-propionamidehydrochloride.

[0591] In a still further embodiment, the serotonin-modulating agent is selected from the group consisting of compounds having the general Formula II shown below and containing, by way of example and not limitation, the compounds listed below. Furthermore, the serotonin-modulating agents useful in the practice of the present invention are described in U.S. Patent No. 5,559,143 which is herein incorporated by reference in its entirety.



Formula II

[0592] wherein:

[0593] B is a C₁₋₄ alkylene bridging group;

[0594] Alk is a linear alkylene bridging group containing from 2-8 carbon atoms which may optionally be mono-substituted at one carbon atom with a C₁₋₄ alkyl, phenyl, substituted phenyl or an alkylphenyl substituent in which the phenyl ring may be optionally substituted;

[0595] D is a bond or an ethylene bridging group;


[0596] X, Y, and Z are each independently hydrogen, C₁₋₄ alkyl, phenyl, substituted phenyl or alkylphenyl in which the phenyl ring may be optionally substituted;

[0597] R₁ is a substituent selected from the group consisting of hydrogen, halogen, C₁₋₄ alkyl, C₁₋₅ alkoxy, CF₃, OCF₃, OH, NO₂, CN, -CONR₂R₃, -NR₂R₃, -COOR₄, -OCH₂COOR₄, -CH₂SO₂NR₂R₃, and -SO₂NR₂R₃;

[0598] R₂ and R₃ are each independently H or a C₁₋₄ alkyl;

[0599] R_4 is H, C_{1-4} alkyl, phenyl, substituted phenyl or an alkylphenyl substituent in which the phenyl ring may be optionally substituted;

[0600] Het is one of the following substituents:



[0601] wherein:

[0602] R is a substituent selected from the group consisting of hydrogen, halogen, C_{1-4} alkyl, C_{1-4} alkoxy, $-O-CH_2-C_6H_5$, CF_3 , OCF_3 , OH, NO_2 , CN, $-CONR_5R_6$, $-CH_2SO_2NR_5R_6$, $-SO_2NR_5R_6$, $-COOR_7$, or $-OCH_2COOR_7$;

[0603] R_5 and R_6 are each independently H or C_{1-4} alkyl;

[0604] R_7 is H, C_{1-4} alkyl, phenyl, substituted phenyl or an alkylphenyl substituent in which the phenyl ring may be optionally substituted; and

[0605] A is H, or C_{1-4} alkyl,

[0606] provided however, that when Het is an indolyl derivative, then R_1 is not a carbonyl derivative.

[0607] Examples of suitable compounds having formula II include:

[0608] 6-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]amino]-N-[(4-(trifluoromethyl)phenyl)]-heptanamide;

[0609] 7-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]amino]-N-(4-methoxyphenyl)-octanamide;

[0610] 6-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]amino]-N-phenylheptanamide;

[0611] 5-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]amino]-N-[4-(trifluoromethyl)phenyl]-hexanamide;

[0612] 6-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]amino]-N-(4-methoxyphenyl)-heptanamide;

[0613] 4-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]amino]-N-[4-(trifluoromethyl)phenyl]-pentanamide;

[0614] 6-[[2-(5-methoxy-1H-indol-3-yl)ethyl]amino]-N-[4-(trifluoromethyl)phenyl]-heptanamide;

[0615] 6-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]methylamino]-N-[4-(trifluoromethyl)phenyl]-heptanamide;

- [0616] 6-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]amino]-N-(2-methoxyphenyl)-heptanamide;
- [0617] 6-[[2-(5-carboxamido-1H-indol-3-yl)ethyl]amino]-N-(4-methoxyphenyl)-heptanamide;
- [0618] 6-[[2-(1H-indol-3-yl)ethyl]amino]-N-[4-(trifluoromethyl)phenyl]-hexanamide;
- [0619] 6-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]amino]-N-[4-(1propyl)phenyl]-hexanamide;
- [0620] 5-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]amino]-N-[4-(1propyloxy)phenyl]-hexanamide;
- [0621] 6-[2-[(2,3-dihydro-1,4-benzodioxin-2-yl)ethyl]amino]-N-phenyl-hexanamide;
- [0622] 6-[(2,3-dihydro-1,4-benzodioxin-2-yl)methylamino]-N-[4-(trifluoromethyl)phenyl]-heptanamide;
- [0623] 6-[(2,3-dihydro-1,4-benzodioxin-2-yl)methylamino]-N-(4-methoxyphenyl)-heptanamide;
- [0624] 6-[(2,3-dihydro-1,4-benzodioxin-2-yl)methyl]-methylamino]-N-[4-(trifluoromethyl)phenyl]-hexanamide;
- [0625] 6-[(2,3-dihydro-1,4-benzodioxin-2-yl)methylamino]-N-[2-(trifluoromethyl)phenyl]-hexanamide;
- [0626] 7-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]amino]-N-(4-methoxyphenyl)-heptanamide;
- [0627] 7-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]amino]-N-(2-methoxyphenyl)-heptanamide;
- [0628] 6-[[2-(4-hydroxy-1H-indol-3-yl)ethyl]amino]-N-(4-methoxyphenyl)-heptanamide;
- [0629] 6-[[2-(5-chloro-1H-indol-3-yl)ethyl]amino]-N-(4-methoxyphenyl)-heptanamide;
- [0630] 7-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]amino]-N-(3-methoxyphenyl)-octanamide;
- [0631] 6-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]amino]-N-[2-(trifluoromethyl)phenyl]-hexanamide;
- [0632] 6-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]amino]-N-[3-(trifluoromethyl)phenyl]-hexanamide;

- [0633] 6-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]amino]-4-methyl-N-(4-methoxyphenyl)-hexanamide;
- [0634] 6-[[3-(5-hydroxy-1H-indol-3-yl)propyl]amino]-N-(4-methoxyphenyl)-hexanamide;
- [0635] 6-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]amino]-N-(3-methoxyphenyl)-hexanamide;
- [0636] 6-[[2-(5-hydroxy-1-methyl-indol-3-yl)ethyl]amino]-N-(4-methoxyphenyl)-hexanamide;
- [0637] 6-[(2,3-dihydro-8-methoxy-1,4-benzodioxin-2-yl)methylamino]-N-(4-methoxyphenyl)-hexanamide;
- [0638] 5-[(2,3-dihydro-1,4-benzodioxin-2-yl)methylamino]-N-[4(trifluoromethyl)phenyl]-pentanamide;
- [0639] 4-[(2,3-dihydro-1,4-benzodioxin-2-yl)methylamino]-N-(4-methoxyphenyl)-butanamide;
- [0640] 7-[[2-(5-methoxy-1H-indol-3-yl)ethyl]-methylamino]-N-(4-methoxyphenyl)-octanamide;
- [0641] 6-[[2-(5-hydroxy-1H-indol-3-yl)ethyl]amino]-N-(4-methoxyphenyl)-2-hexenamide; and
- [0642] 7-[(2,3-dihydro-1,4-benzodioxin-2-yl)methylamino]-N-[4-trifluoromethyl)phenyl]-heptanamide.

[0643] In yet another aspect of the invention, the serotonin-modulating agent is a serotonin reuptake inhibitor. In one embodiment, the serotonin reuptake inhibitor is citalopram (marketed under the trademark Celexa[®] by Forest Laboratories, Parke-Davis, Inc). In another embodiment, the serotonin reuptake inhibitor is fluoxetine (marketed under the trademark Prozac[®] by Eli Lilly and Company). In still another embodiment, the serotonin reuptake inhibitor is fluvoxamine (marketed under the trademark Luvox[®] by Solvay Pharmaceuticals, Inc.). In yet another embodiment, the serotonin reuptake inhibitor is paroxetine (marketed under the trademark Paxil[®] by SmithKline Beecham Pharmaceuticals, Inc.). In a further embodiment, the serotonin reuptake inhibitor is escitalopram oxalate (marketed under the trademark Lexapro[®] by Forest Laboratories, Parke-Davis, Inc). In still another embodiment, the serotonin reuptake inhibitor is sertraline (marketed under the trademark Zoloft[®] by Pfizer, Inc.).

[0644] It is also contemplated that a metabolite of a serotonin reuptake inhibitor may be used in the practice of the invention. Any metabolite of a serotonin reuptake inhibitor may be employed in the current invention to the extent that the compound reverse

or lessen central nervous system cell damage following a reduction in blood flow to the central nervous system. By way of example, in one embodiment, the metabolite is norfluoxetine, which is an active metabolite of fluoxetine. By way of further example, in another embodiment, the metabolite is N-demethylsertraline, which is an active metabolite of sertraline.

[0645] Generally speaking, the pharmacokinetics of the particular agent to be administered will dictate the most preferred method of administration and dosing regimen. The serotonin-modulating agent can be administered as a pharmaceutical composition with or without a carrier. The terms "pharmaceutically acceptable carrier" or a "carrier" refer to any generally acceptable excipient or drug delivery composition that is relatively inert and non-toxic. Exemplary carriers include sterile water, salt solutions (such as Ringer's solution), alcohols, gelatin, talc, viscous paraffin, fatty acid esters, hydroxymethylcellulose, polyvinyl pyrrolidone, calcium carbonate, carbohydrates (such as lactose, sucrose, dextrose, and mannose), albumin, starch, cellulose, silica gel, polyethylene glycol (PEG), dried skim milk, rice flour, magnesium stearate, and the like. Suitable formulations and additional carriers are described in Remington's Pharmaceutical Sciences, (17th Ed., Mack Pub. Co., Easton, Pa.). Such preparations can be sterilized and, if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, preservatives and/or aromatic substances and the like which do not deleteriously react with the active compounds. Typical preservatives can include, potassium sorbate, sodium metabisulfite, methyl paraben, propyl paraben, thimerosal, etc. The compositions can also be combined where desired with other active substances, e.g., enzyme inhibitors, to reduce metabolic degradation.

[0646] Moreover, the serotonin-modulating agent can be a liquid solution, suspension, emulsion, tablet, pill, capsule, sustained release formulation, or powder. The method of administration can dictate how the composition will be formulated. For example, the composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulations can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, cellulose, or magnesium carbonate.

[0647] In another embodiment, the serotonin-modulating agent can be administered intravenously, parenterally, intramuscular, subcutaneously, orally, nasally, topically, by inhalation, by implant, by injection, or by suppository. For enteral or mucosal application (including via oral and nasal mucosa), particularly suitable are tablets, liquids,

drops, suppositories or capsules. A syrup, elixir or the like can be used wherein a sweetened vehicle is employed. Liposomes, microspheres, and microcapsules are available and can be used. Pulmonary administration can be accomplished, for example, using any of various delivery devices known in the art such as an inhaler. See, e.g. S. P. Newman (1984) in *Aerosols and the Lung*, Clarke and Davis (eds.), Butterworths, London, England, pp. 197-224; PCT Publication No. WO 92/16192; PCT Publication No. WO 91/08760. For parenteral application, particularly suitable are injectable, sterile solutions, preferably oily or aqueous solutions, as well as suspensions, emulsions, or implants. In particular, carriers for parenteral administration include aqueous solutions of dextrose, saline, pure water, ethanol, glycerol, propylene glycol, peanut oil, sesame oil, polyoxyethylene-polyoxypropylene block polymers, and the like.

[0648] The actual effective amounts of compound or drug can and will vary according to the specific composition being utilized, the mode of administration and the age, weight and condition of the subject. Dosages for a particular individual subject can be determined by one of ordinary skill in the art using conventional considerations. But in general, the amount of serotonin-modulating agent will be between about 10 to about 2500 milligrams per day. The daily dose can be administered in one to four doses per day.

[0649] In one embodiment, when the serotonin-modulating agent comprises sertraline, typically the amount administered is within a range of from about 0.5 to about 200 milligrams per day, and even more typically, between about 50 to about 100 milligrams per day.

[0650] In another embodiment, when the serotonin-modulating agent is fluvoxamine, typically the amount administered is within a range of from about 0.5 to about 500 milligrams per day, and even more typically, between about 100 to about 300 milligrams per day.

[0651] In yet another embodiment, when the serotonin-modulating agent is fluoxetine, generally the amount administered is within a range of from about 0.5 to about 150 milligrams per day, and even more typically, between about 20 to about 80 milligrams per day.

[0652] In still another embodiment, when the serotonin-modulating agent is paroxetine, typically the amount administered is within a range of from about 0.5 to about 100 milligrams per day, and even more typically, between about 10 to about 50 milligrams per day.

[0653] In yet a further embodiment, when the serotonin-modulating agent is citalopram, typically the amount administered is within a range of from about 0.5 to about

100 milligrams per day, and even more typically, between about 20 to about 40 milligrams per day.

[0654] In still another embodiment, when the serotonin-modulating agent is escitalopram oxalate, typically the amount administered is within a range of from about 0.5 to about 50 milligrams per day, and even more typically, between about 5 to about 20 milligrams per day.

[0655] Generally speaking, when the composition is administered to treat an ischemic mediated condition, the serotonin-modulating agent and cyclooxygenase-2 selective inhibitor are administered to the subject as soon as possible after the reduction in blood flow to the central nervous system in order to reduce the extent of ischemic damage. Typically, the serotonin-modulating agent and cyclooxygenase-2 selective inhibitor are administered within 10 days after the reduction of blood flow to the central nervous system and more typically, within 24 hours. In still another embodiment, the serotonin-modulating agent and cyclooxygenase-2 selective inhibitor are administered from about 1 to about 12 hours after the reduction in blood flow to the central nervous system. In another embodiment, the serotonin-modulating agent and cyclooxygenase-2 selective inhibitor are administered in less than about 6 hours after the reduction in blood flow to the central nervous system. In still another embodiment, the serotonin-modulating agent and cyclooxygenase-2 selective inhibitor are administered in less than about 4 hours after the reduction in blood flow to the central nervous system. In yet a further embodiment, the serotonin-modulating agent and cyclooxygenase-2 selective inhibitor are administered in less than about 2 hours after the reduction in blood flow to the central nervous system.

[0656] Moreover, the timing of the administration of the cyclooxygenase-2 selective inhibitor in relation to the administration of the serotonin-modulating agent may also vary from subject to subject. In one embodiment, the cyclooxygenase-2 selective inhibitor and serotonin-modulating agent may be administered substantially simultaneously, meaning that both agents may be administered to the subject at approximately the same time. For example, the cyclooxygenase-2 selective is administered during a continuous period beginning on the same day as the beginning of the serotonin-modulating agent and extending to a period after the end of the serotonin-modulating agent. Alternatively, the cyclooxygenase-2 selective inhibitor and serotonin-modulating agent may be administered sequentially, meaning that they are administered at separate times during separate treatments. In one embodiment, for example, the cyclooxygenase-2 selective inhibitor is administered during a continuous period beginning prior to administration of the serotonin-modulating agent and ending after administration of

the serotonin-modulating agent. Of course, it is also possible that the cyclooxygenase-2 selective inhibitor may be administered either more or less frequently than the serotonin-modulating agent. Moreover, it will be apparent to those skilled in the art that it is possible, and perhaps desirable, to combine various times and methods of administration in the practice of the present invention.

COMBINATION THERAPIES

[0657] Generally speaking, it is contemplated that the composition employed in the practice of the invention may include one or more of any of the cyclooxygenase-2 selective inhibitors detailed above in combination with one or more of any of the serotonin-modulating agents detailed above. By way of a non-limiting example, Table 4a details a number of suitable combinations that are useful in the methods and compositions of the current invention. The combination may also include an isomer, a pharmaceutically acceptable salt, ester, or prodrug of any of the cyclooxygenase-2 selective inhibitors or serotonin-modulating agents listed in Table 4a.

TABLE 4a

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
a compound having formula I	citalopram
a compound having formula I	fluoxetine
a compound having formula I	fluvoxamine
a compound having formula I	paroxetine
a compound having formula I	6-chloro-2-[piperidinyl-4-thio]pyridine
a compound having formula I	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
a compound having formula I	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
a compound having formula I	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
a compound having formula I	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
a compound having formula I	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
a compound having formula II	citalopram
a compound having formula II	fluoxetine
a compound having formula II	fluvoxamine
a compound having formula II	paroxetine

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
a compound having formula II	6-chloro-2-[piperidinyl-4-thio]pyridine
a compound having formula II	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
a compound having formula II	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
a compound having formula II	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
a compound having formula II	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
a compound having formula II	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
a compound having formula III	citalopram
a compound having formula III	fluoxetine
a compound having formula III	fluvoxamine
a compound having formula III	paroxetine
a compound having formula III	6-chloro-2-[piperidinyl-4-thio]pyridine
a compound having formula III	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
a compound having formula III	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
a compound having formula III	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
a compound having formula III	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
a compound having formula III	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
a compound having formula IV	citalopram
a compound having formula IV	fluoxetine
a compound having formula IV	fluvoxamine
a compound having formula IV	paroxetine
a compound having formula IV	6-chloro-2-[piperidinyl-4-thio]pyridine
a compound having formula IV	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
a compound having formula IV	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
a compound having formula IV	3-[4-(4-chlorophenyl)piperazin-1-

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
	yl]-1,1-diphenyl-2-propanol
a compound having formula IV	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
a compound having formula IV	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
a compound having formula V	citalopram
a compound having formula V	fluoxetine
a compound having formula V	fluvoxamine
a compound having formula V	paroxetine
a compound having formula V	6-chloro-2-[piperidinyl-4-thio]pyridine
a compound having formula V	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
a compound having formula V	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
a compound having formula V	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
a compound having formula V	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
a compound having formula V	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide

[0658] By way of further example, Table 4b details a number of suitable combinations that may be employed in the methods and compositions of the present invention. The combination may also include an isomer, a pharmaceutically acceptable salt, ester, or prodrug of any of the cyclooxygenase-2 selective inhibitors or serotonin-modulating agents listed in Table 4b.

TABLE 4b

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
a compound selected from the group consisting of B-1, B-2, B-3, B-4, B-5, B-6, B-7, B-8, B-9, B-10, B-11, B-12, B-13, B-14, B-15, B-16, B-17, B-18, B-19, B-20, B-21, B-22, B-23, B23a, B-24, B-25, B-26, B-27, B-28, B-29, B-30, B-31, B-32, B-33, B-34, B-35, B-36, B-37, B-38, B-39, B-40, B-41, B-42, B-43, B-44, B-45, B-46, B-47, B-48, B-49, B-50, B-51, B-52, B-53, B-54, B-55, B-56, B-57, B-58, B-59, B-60, B-61, B-62, B-63, B-64, B-65, B-66, B-67, B-68, B-69, B-70, B-71, B-72, B-73, B-74, B-75, B-76, B-77, B-78, B-79, B-80, B-81, B-82, B-83, B-84, B-85, B-86, B-87, B-88, B-89, B-90, B-91, B-92, B-93, B-94, B-95, B-96, B-97, B-98, B-99, B-100, B-101, B-102, B-103, B-104, B-105, B-106, B-107, B-108, B-109, B-110, B-111, B-112, B-113, B-114, B-115, B-116, B-117, B-118, B-119, B-120, B-121, B-122, B-123, B-124, B-125, B-126, B-127, B-128, B-129, B-130, B-131, B-132, B-133, B-134, B-135, B-136, B-137, B-138, B-139, B-140, B-141, B-142, B-143, B-144, B-145, B-146, B-147, B-148, B-149, B-150, B-151, B-152, B-153, B-154, B-155, B-156, B-157, B-158, B-159, B-160, B-161, B-162, B-163, B-164, B-165, B-166, B-167, B-168, B-169, B-170, B-171, B-172, B-173, B-174, B-175, B-176, B-177, B-178, B-179, B-180, B-181, B-182, B-183, B-184, B-185, B-186, B-187, B-188, B-189, B-190, B-191, B-192, B-193, B-194, B-195, B-196, B-197, B-198, B-199, B-200, B-201, B-202, B-203, B-204, B-205, B-206, B-207, B-208, B-209, B-210, B-211, B-212, B-213, B-214, B-215, B-216, B-217, B-218, B-219, B-220, B-221, B-222, B-223, B-224, B-225, B-226, B-227, B-228, B-229, B-230, B-231, B-232, B233, B-234, B-235, B-236, B-237, B-238, B-239, B-240, B-241, B-242, B-243, B-244, B-245, B-246, B-247, B-248, B-249, B-250, B-251, and B-252.	citalopram

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
<p>a compound selected from the group consisting of B-1, B-2, B-3, B-4, B-5, B-6, B-7, B-8, B-9, B-10, B-11, B-12, B-13, B-14, B-15, B-16, B-17, B-18, B-19, B-20, B-21, B-22, B-23, B23a, B-24, B-25, B-26, B-27, B-28, B-29, B-30, B-31, B-32, B-33, B-34, B-35, B-36, B-37, B-38, B-39, B-40, B-41, B-42, B-43, B-44, B-45, B-46, B-47, B-48, B-49, B-50, B-51, B-52, B-53, B-54, B-55, B-56, B-57, B-58, B-59, B-60, B-61, B-62, B-63, B-64, B-65, B-66, B-67, B-68, B-69, B-70, B-71, B-72, B-73, B-74, B-75, B-76, B-77, B-78, B-79, B-80, B-81, B-82, B-83, B-84, B-85, B-86, B-87, B-88, B-89, B-90, B-91, B-92, B-93, B-94, B-95, B-96, B-97, B-98, B-99, B-100, B-101, B-102, B-103, B-104, B-105, B-106, B-107, B-108, B-109, B-110, B-111, B-112, B-113, B-114, B-115, B-116, B-117, B-118, B-119, B-120, B-121, B-122, B-123, B-124, B-125, B-126, B-127, B-128, B-129, B-130, B-131, B-132, B-133, B-134, B-135, B-136, B-137, B-138, B-139, B-140, B-141, B-142, B-143, B-144, B-145, B-146, B-147, B-148, B-149, B-150, B-151, B-152, B-153, B-154, B-155, B-156, B-157, B-158, B-159, B-160, B-161, B-162, B-163, B-164, B-165, B-166, B-167, B-168, B-169, B-170, B-171, B-172, B-173, B-174, B-175, B-176, B-177, B-178, B-179, B-180, B-181, B-182, B-183, B-184, B-185, B-186, B-187, B-188, B-189, B-190, B-191, B-192, B-193, B-194, B-195, B-196, B-197, B-198, B-199, B-200, B-201, B-202, B-203, B-204, B-205, B-206, B-207, B-208, B-209, B-210, B-211, B-212, B-213, B-214, B-215, B-216, B-217, B-218, B-219, B-220, B-221, B-222, B-223, B-224, B-225, B-226, B-227, B-228, B-229, B-230, B-231, B-232, B233, B-234, B-235, B-236, B-237, B-238, B-239, B-240, B-241, B-242, B-243, B-244, B-245, B-246, B-247, B-248, B-249, B-250, B-251, and B-252.</p>	<p>fluoxetine</p>

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
a compound selected from the group consisting of B-1, B-2, B-3, B-4, B-5, B-6, B-7, B-8, B-9, B-10, B-11, B-12, B-13, B-14, B-15, B-16, B-17, B-18, B-19, B-20, B-21, B-22, B-23, B23a, B-24, B-25, B-26, B-27, B-28, B-29, B-30, B-31, B-32, B-33, B-34, B-35, B-36, B-37, B-38, B-39, B-40, B-41, B-42, B-43, B-44, B-45, B-46, B-47, B-48, B-49, B-50, B-51, B-52, B-53, B-54, B-55, B-56, B-57, B-58, B-59, B-60, B-61, B-62, B-63, B-64, B-65, B-66, B-67, B-68, B-69, B-70, B-71, B-72, B-73, B-74, B-75, B-76, B-77, B-78, B-79, B-80, B-81, B-82, B-83, B-84, B-85, B-86, B-87, B-88, B-89, B-90, B-91, B-92, B-93, B-94, B-95, B-96, B-97, B-98, B-99, B-100, B-101, B-102, B-103, B-104, B-105, B-106, B-107, B-108, B-109, B-110, B-111, B-112, B-113, B-114, B-115, B-116, B-117, B-118, B-119, B-120, B-121, B-122, B-123, B-124, B-125, B-126, B-127, B-128, B-129, B-130, B-131, B-132, B-133, B-134, B-135, B-136, B-137, B-138, B-139, B-140, B-141, B-142, B-143, B-144, B-145, B-146, B-147, B-148, B-149, B-150, B-151, B-152, B-153, B-154, B-155, B-156, B-157, B-158, B-159, B-160, B-161, B-162, B-163, B-164, B-165, B-166, B-167, B-168, B-169, B-170, B-171, B-172, B-173, B-174, B-175, B-176, B-177, B-178, B-179, B-180, B-181, B-182, B-183, B-184, B-185, B-186, B-187, B-188, B-189, B-190, B-191, B-192, B-193, B-194, B-195, B-196, B-197, B-198, B-199, B-200, B-201, B-202, B-203, B-204, B-205, B-206, B-207, B-208, B-209, B-210, B-211, B-212, B-213, B-214, B-215, B-216, B-217, B-218, B-219, B-220, B-221, B-222, B-223, B-224, B-225, B-226, B-227, B-228, B-229, B-230, B-231, B-232, B233, B-234, B-235, B-236, B-237, B-238, B-239, B-240, B-241, B-242, B-243, B-244, B-245, B-246, B-247, B-248, B-249, B-250, B-251, and B-252.	fluvoxamine

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
a compound selected from the group consisting of B-1, B-2, B-3, B-4, B-5, B-6, B-7, B-8, B-9, B-10, B-11, B-12, B-13, B-14, B-15, B-16, B-17, B-18, B-19, B-20, B-21, B-22, B-23, B23a, B-24, B-25, B-26, B-27, B-28, B-29, B-30, B-31, B-32, B-33, B-34, B-35, B-36, B-37, B-38, B-39, B-40, B-41, B-42, B-43, B-44, B-45, B-46, B-47, B-48, B-49, B-50, B-51, B-52, B-53, B-54, B-55, B-56, B-57, B-58, B-59, B-60, B-61, B-62, B-63, B-64, B-65, B-66, B-67, B-68, B-69, B-70, B-71, B-72, B-73, B-74, B-75, B-76, B-77, B-78, B-79, B-80, B-81, B-82, B-83, B-84, B-85, B-86, B-87, B-88, B-89, B-90, B-91, B-92, B-93, B-94, B-95, B-96, B-97, B-98, B-99, B-100, B-101, B-102, B-103, B-104, B-105, B-106, B-107, B-108, B-109, B-110, B-111, B-112, B-113, B-114, B-115, B-116, B-117, B-118, B-119, B-120, B-121, B-122, B-123, B-124, B-125, B-126, B-127, B-128, B-129, B-130, B-131, B-132, B-133, B-134, B-135, B-136, B-137, B-138, B-139, B-140, B-141, B-142, B-143, B-144, B-145, B-146, B-147, B-148, B-149, B-150, B-151, B-152, B-153, B-154, B-155, B-156, B-157, B-158, B-159, B-160, B-161, B-162, B-163, B-164, B-165, B-166, B-167, B-168, B-169, B-170, B-171, B-172, B-173, B-174, B-175, B-176, B-177, B-178, B-179, B-180, B-181, B-182, B-183, B-184, B-185, B-186, B-187, B-188, B-189, B-190, B-191, B-192, B-193, B-194, B-195, B-196, B-197, B-198, B-199, B-200, B-201, B-202, B-203, B-204, B-205, B-206, B-207, B-208, B-209, B-210, B-211, B-212, B-213, B-214, B-215, B-216, B-217, B-218, B-219, B-220, B-221, B-222, B-223, B-224, B-225, B-226, B-227, B-228, B-229, B-230, B-231, B-232, B233, B-234, B-235, B-236, B-237, B-238, B-239, B-240, B-241, B-242, B-243, B-244, B-245, B-246, B-247, B-248, B-249, B-250, B-251, and B-252.	paroxetine

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
<p>a compound selected from the group consisting of B-1, B-2, B-3, B-4, B-5, B-6, B-7, B-8, B-9, B-10, B-11, B-12, B-13, B-14, B-15, B-16, B-17, B-18, B-19, B-20, B-21, B-22, B-23, B23a, B-24, B-25, B-26, B-27, B-28, B-29, B-30, B-31, B-32, B-33, B-34, B-35, B-36, B-37, B-38, B-39, B-40, B-41, B-42, B-43, B-44, B-45, B-46, B-47, B-48, B-49, B-50, B-51, B-52, B-53, B-54, B-55, B-56, B-57, B-58, B-59, B-60, B-61, B-62, B-63, B-64, B-65, B-66, B-67, B-68, B-69, B-70, B-71, B-72, B-73, B-74, B-75, B-76, B-77, B-78, B-79, B-80, B-81, B-82, B-83, B-84, B-85, B-86, B-87, B-88, B-89, B-90, B-91, B-92, B-93, B-94, B-95, B-96, B-97, B-98, B-99, B-100, B-101, B-102, B-103, B-104, B-105, B-106, B-107, B-108, B-109, B-110, B-111, B-112, B-113, B-114, B-115, B-116, B-117, B-118, B-119, B-120, B-121, B-122, B-123, B-124, B-125, B-126, B-127, B-128, B-129, B-130, B-131, B-132, B-133, B-134, B-135, B-136, B-137, B-138, B-139, B-140, B-141, B-142, B-143, B-144, B-145, B-146, B-147, B-148, B-149, B-150, B-151, B-152, B-153, B-154, B-155, B-156, B-157, B-158, B-159, B-160, B-161, B-162, B-163, B-164, B-165, B-166, B-167, B-168, B-169, B-170, B-171, B-172, B-173, B-174, B-175, B-176, B-177, B-178, B-179, B-180, B-181, B-182, B-183, B-184, B-185, B-186, B-187, B-188, B-189, B-190, B-191, B-192, B-193, B-194, B-195, B-196, B-197, B-198, B-199, B-200, B-201, B-202, B-203, B-204, B-205, B-206, B-207, B-208, B-209, B-210, B-211, B-212, B-213, B-214, B-215, B-216, B-217, B-218, B-219, B-220, B-221, B-222, B-223, B-224, B-225, B-226, B-227, B-228, B-229, B-230, B-231, B-232, B233, B-234, B-235, B-236, B-237, B-238, B-239, B-240, B-241, B-242, B-243, B-244, B-245, B-246, B-247, B-248, B-249, B-250, B-251, and B-252.</p>	<p>6-chloro-2-[piperidinyl-4-thio]pyridine</p>

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
<p>a compound selected from the group consisting of B-1, B-2, B-3, B-4, B-5, B-6, B-7, B-8, B-9, B-10, B-11, B-12, B-13, B-14, B-15, B-16, B-17, B-18, B-19, B-20, B-21, B-22, B-23, B23a, B-24, B-25, B-26, B-27, B-28, B-29, B-30, B-31, B-32, B-33, B-34, B-35, B-36, B-37, B-38, B-39, B-40, B-41, B-42, B-43, B-44, B-45, B-46, B-47, B-48, B-49, B-50, B-51, B-52, B-53, B-54, B-55, B-56, B-57, B-58, B-59, B-60, B-61, B-62, B-63, B-64, B-65, B-66, B-67, B-68, B-69, B-70, B-71, B-72, B-73, B-74, B-75, B-76, B-77, B-78, B-79, B-80, B-81, B-82, B-83, B-84, B-85, B-86, B-87, B-88, B-89, B-90, B-91, B-92, B-93, B-94, B-95, B-96, B-97, B-98, B-99, B-100, B-101, B-102, B-103, B-104, B-105, B-106, B-107, B-108, B-109, B-110, B-111, B-112, B-113, B-114, B-115, B-116, B-117, B-118, B-119, B-120, B-121, B-122, B-123, B-124, B-125, B-126, B-127, B-128, B-129, B-130, B-131, B-132, B-133, B-134, B-135, B-136, B-137, B-138, B-139, B-140, B-141, B-142, B-143, B-144, B-145, B-146, B-147, B-148, B-149, B-150, B-151, B-152, B-153, B-154, B-155, B-156, B-157, B-158, B-159, B-160, B-161, B-162, B-163, B-164, B-165, B-166, B-167, B-168, B-169, B-170, B-171, B-172, B-173, B-174, B-175, B-176, B-177, B-178, B-179, B-180, B-181, B-182, B-183, B-184, B-185, B-186, B-187, B-188, B-189, B-190, B-191, B-192, B-193, B-194, B-195, B-196, B-197, B-198, B-199, B-200, B-201, B-202, B-203, B-204, B-205, B-206, B-207, B-208, B-209, B-210, B-211, B-212, B-213, B-214, B-215, B-216, B-217, B-218, B-219, B-220, B-221, B-222, B-223, B-224, B-225, B-226, B-227, B-228, B-229, B-230, B-231, B-232, B233, B-234, B-235, B-236, B-237, B-238, B-239, B-240, B-241, B-242, B-243, B-244, B-245, B-246, B-247, B-248, B-249, B-250, B-251, and B-252.</p>	<p>8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione</p>

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
<p>a compound selected from the group consisting of B-1, B-2, B-3, B-4, B-5, B-6, B-7, B-8, B-9, B-10, B-11, B-12, B-13, B-14, B-15, B-16, B-17, B-18, B-19, B-20, B-21, B-22, B-23, B23a, B-24, B-25, B-26, B-27, B-28, B-29, B-30, B-31, B-32, B-33, B-34, B-35, B-36, B-37, B-38, B-39, B-40, B-41, B-42, B-43, B-44, B-45, B-46, B-47, B-48, B-49, B-50, B-51, B-52, B-53, B-54, B-55, B-56, B-57, B-58, B-59, B-60, B-61, B-62, B-63, B-64, B-65, B-66, B-67, B-68, B-69, B-70, B-71, B-72, B-73, B-74, B-75, B-76, B-77, B-78, B-79, B-80, B-81, B-82, B-83, B-84, B-85, B-86, B-87, B-88, B-89, B-90, B-91, B-92, B-93, B-94, B-95, B-96, B-97, B-98, B-99, B-100, B-101, B-102, B-103, B-104, B-105, B-106, B-107, B-108, B-109, B-110, B-111, B-112, B-113, B-114, B-115, B-116, B-117, B-118, B-119, B-120, B-121, B-122, B-123, B-124, B-125, B-126, B-127, B-128, B-129, B-130, B-131, B-132, B-133, B-134, B-135, B-136, B-137, B-138, B-139, B-140, B-141, B-142, B-143, B-144, B-145, B-146, B-147, B-148, B-149, B-150, B-151, B-152, B-153, B-154, B-155, B-156, B-157, B-158, B-159, B-160, B-161, B-162, B-163, B-164, B-165, B-166, B-167, B-168, B-169, B-170, B-171, B-172, B-173, B-174, B-175, B-176, B-177, B-178, B-179, B-180, B-181, B-182, B-183, B-184, B-185, B-186, B-187, B-188, B-189, B-190, B-191, B-192, B-193, B-194, B-195, B-196, B-197, B-198, B-199, B-200, B-201, B-202, B-203, B-204, B-205, B-206, B-207, B-208, B-209, B-210, B-211, B-212, B-213, B-214, B-215, B-216, B-217, B-218, B-219, B-220, B-221, B-222, B-223, B-224, B-225, B-226, B-227, B-228, B-229, B-230, B-231, B-232, B233, B-234, B-235, B-236, B-237, B-238, B-239, B-240, B-241, B-242, B-243, B-244, B-245, B-246, B-247, B-248, B-249, B-250, B-251, and B-252.</p>	<p>1-[3-(3,4-methylenedioxy phenoxy)propyl]-4-phenylpiperazine</p>

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
<p>a compound selected from the group consisting of B-1, B-2, B-3, B-4, B-5, B-6, B-7, B-8, B-9, B-10, B-11, B-12, B-13, B-14, B-15, B-16, B-17, B-18, B-19, B-20, B-21, B-22, B-23, B23a, B-24, B-25, B-26, B-27, B-28, B-29, B-30, B-31, B-32, B-33, B-34, B-35, B-36, B-37, B-38, B-39, B-40, B-41, B-42, B-43, B-44, B-45, B-46, B-47, B-48, B-49, B-50, B-51, B-52, B-53, B-54, B-55, B-56, B-57, B-58, B-59, B-60, B-61, B-62, B-63, B-64, B-65, B-66, B-67, B-68, B-69, B-70, B-71, B-72, B-73, B-74, B-75, B-76, B-77, B-78, B-79, B-80, B-81, B-82, B-83, B-84, B-85, B-86, B-87, B-88, B-89, B-90, B-91, B-92, B-93, B-94, B-95, B-96, B-97, B-98, B-99, B-100, B-101, B-102, B-103, B-104, B-105, B-106, B-107, B-108, B-109, B-110, B-111, B-112, B-113, B-114, B-115, B-116, B-117, B-118, B-119, B-120, B-121, B-122, B-123, B-124, B-125, B-126, B-127, B-128, B-129, B-130, B-131, B-132, B-133, B-134, B-135, B-136, B-137, B-138, B-139, B-140, B-141, B-142, B-143, B-144, B-145, B-146, B-147, B-148, B-149, B-150, B-151, B-152, B-153, B-154, B-155, B-156, B-157, B-158, B-159, B-160, B-161, B-162, B-163, B-164, B-165, B-166, B-167, B-168, B-169, B-170, B-171, B-172, B-173, B-174, B-175, B-176, B-177, B-178, B-179, B-180, B-181, B-182, B-183, B-184, B-185, B-186, B-187, B-188, B-189, B-190, B-191, B-192, B-193, B-194, B-195, B-196, B-197, B-198, B-199, B-200, B-201, B-202, B-203, B-204, B-205, B-206, B-207, B-208, B-209, B-210, B-211, B-212, B-213, B-214, B-215, B-216, B-217, B-218, B-219, B-220, B-221, B-222, B-223, B-224, B-225, B-226, B-227, B-228, B-229, B-230, B-231, B-232, B233, B-234, B-235, B-236, B-237, B-238, B-239, B-240, B-241, B-242, B-243, B-244, B-245, B-246, B-247, B-248, B-249, B-250, B-251, and B-252.</p>	<p>3-[4-(4-chlorophenyl) piperazin-1-yl]-1,1-diphenyl-2-propanol</p>

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
<p>a compound selected from the group consisting of B-1, B-2, B-3, B-4, B-5, B-6, B-7, B-8, B-9, B-10, B-11, B-12, B-13, B-14, B-15, B-16, B-17, B-18, B-19, B-20, B-21, B-22, B-23, B23a, B-24, B-25, B-26, B-27, B-28, B-29, B-30, B-31, B-32, B-33, B-34, B-35, B-36, B-37, B-38, B-39, B-40, B-41, B-42, B-43, B-44, B-45, B-46, B-47, B-48, B-49, B-50, B-51, B-52, B-53, B-54, B-55, B-56, B-57, B-58, B-59, B-60, B-61, B-62, B-63, B-64, B-65, B-66, B-67, B-68, B-69, B-70, B-71, B-72, B-73, B-74, B-75, B-76, B-77, B-78, B-79, B-80, B-81, B-82, B-83, B-84, B-85, B-86, B-87, B-88, B-89, B-90, B-91, B-92, B-93, B-94, B-95, B-96, B-97, B-98, B-99, B-100, B-101, B-102, B-103, B-104, B-105, B-106, B-107, B-108, B-109, B-110, B-111, B-112, B-113, B-114, B-115, B-116, B-117, B-118, B-119, B-120, B-121, B-122, B-123, B-124, B-125, B-126, B-127, B-128, B-129, B-130, B-131, B-132, B-133, B-134, B-135, B-136, B-137, B-138, B-139, B-140, B-141, B-142, B-143, B-144, B-145, B-146, B-147, B-148, B-149, B-150, B-151, B-152, B-153, B-154, B-155, B-156, B-157, B-158, B-159, B-160, B-161, B-162, B-163, B-164, B-165, B-166, B-167, B-168, B-169, B-170, B-171, B-172, B-173, B-174, B-175, B-176, B-177, B-178, B-179, B-180, B-181, B-182, B-183, B-184, B-185, B-186, B-187, B-188, B-189, B-190, B-191, B-192, B-193, B-194, B-195, B-196, B-197, B-198, B-199, B-200, B-201, B-202, B-203, B-204, B-205, B-206, B-207, B-208, B-209, B-210, B-211, B-212, B-213, B-214, B-215, B-216, B-217, B-218, B-219, B-220, B-221, B-222, B-223, B-224, B-225, B-226, B-227, B-228, B-229, B-230, B-231, B-232, B233, B-234, B-235, B-236, B-237, B-238, B-239, B-240, B-241, B-242, B-243 B-244, B-245, B-246, B-247, B-248, B-249, B-250, B-251, and B-252.</p>	<p>4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile</p>

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
a compound selected from the group consisting of B-1, B-2, B-3, B-4, B-5, B-6, B-7, B-8, B-9, B-10, B-11, B-12, B-13, B-14, B-15, B-16, B-17, B-18, B-19, B-20, B-21, B-22, B-23, B23a, B-24, B-25, B-26, B-27, B-28, B-29, B-30, B-31, B-32, B-33, B-34, B-35, B-36, B-37, B-38, B-39, B-40, B-41, B-42, B-43, B-44, B-45, B-46, B-47, B-48, B-49, B-50, B-51, B-52, B-53, B-54, B-55, B-56, B-57, B-58, B-59, B-60, B-61, B-62, B-63, B-64, B-65, B-66, B-67, B-68, B-69, B-70, B-71, B-72, B-73, B-74, B-75, B-76, B-77, B-78, B-79, B-80, B-81, B-82, B-83, B-84, B-85, B-86, B-87, B-88, B-89, B-90, B-91, B-92, B-93, B-94, B-95, B-96, B-97, B-98, B-99, B-100, B-101, B-102, B-103, B-104, B-105, B-106, B-107, B-108, B-109, B-110, B-111, B-112, B-113, B-114, B-115, B-116, B-117, B-118, B-119, B-120, B-121, B-122, B-123, B-124, B-125, B-126, B-127, B-128, B-129, B-130, B-131, B-132, B-133, B-134, B-135, B-136, B-137, B-138, B-139, B-140, B-141, B-142, B-143, B-144, B-145, B-146, B-147, B-148, B-149, B-150, B-151, B-152, B-153, B-154, B-155, B-156, B-157, B-158, B-159, B-160, B-161, B-162, B-163, B-164, B-165, B-166, B-167, B-168, B-169, B-170, B-171, B-172, B-173, B-174, B-175, B-176, B-177, B-178, B-179, B-180, B-181, B-182, B-183, B-184, B-185, B-186, B-187, B-188, B-189, B-190, B-191, B-192, B-193, B-194, B-195, B-196, B-197, B-198, B-199, B-200, B-201, B-202, B-203, B-204, B-205, B-206, B-207, B-208, B-209, B-210, B-211, B-212, B-213, B-214, B-215, B-216, B-217, B-218, B-219, B-220, B-221, B-222, B-223, B-224, B-225, B-226, B-227, B-228, B-229, B-230, B-231, B-232, B233, B-234, B-235, B-236, B-237, B-238, B-239, B-240, B-241, B-242, B-243, B-244, B-245, B-246, B-247, B-248, B-249, B-250, B-251, and B-252.	3-[3-(dimethylamino) propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide

[0659] By way of yet further example, Table 4c details additional suitable combinations that may be employed in the methods and compositions of the current invention. The combination may also include an isomer, a pharmaceutically acceptable salt, ester, or prodrug of any of the cyclooxygenase-2 selective inhibitors or serotonin-modulating agents listed in Table 4c.

TABLE 4c

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
celecoxib	citalopram
celecoxib	fluoxetine
celecoxib	fluvoxamine
celecoxib	paroxetine
celecoxib	6-chloro-2-[piperidinyl-4-thio]pyridine
celecoxib	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
celecoxib	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
celecoxib	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
celecoxib	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
celecoxib	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
cimicoxib	citalopram
cimicoxib	fluoxetine
cimicoxib	fluvoxamine
cimicoxib	paroxetine
cimicoxib	6-chloro-2-[piperidinyl-4-thio]pyridine
cimicoxib	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
cimicoxib	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
cimicoxib	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
cimicoxib	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
cimicoxib	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
deracoxib	citalopram
deracoxib	fluoxetine
deracoxib	fluvoxamine
deracoxib	paroxetine
deracoxib	6-chloro-2-[piperidinyl-4-thio]pyridine
deracoxib	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
deracoxib	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
deracoxib	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
deracoxib	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
deracoxib	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
valdecoxib	citalopram
valdecoxib	fluoxetine
valdecoxib	fluvoxamine
valdecoxib	paroxetine
valdecoxib	6-chloro-2-[piperidinyl-4-thio]pyridine
valdecoxib	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
valdecoxib	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
valdecoxib	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
valdecoxib	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
valdecoxib	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
rofecoxib	citalopram
rofecoxib	fluoxetine
rofecoxib	fluvoxamine
rofecoxib	paroxetine
rofecoxib	6-chloro-2-[piperidinyl-4-thio]pyridine
rofecoxib	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
rofecoxib	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
rofecoxib	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
rofecoxib	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
rofecoxib	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
etoricoxib	citalopram
etoricoxib	fluoxetine
etoricoxib	fluvoxamine
etoricoxib	paroxetine
etoricoxib	6-chloro-2-[piperidinyl-4-thio]pyridine
etoricoxib	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
etoricoxib	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
etoricoxib	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
etoricoxib	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
etoricoxib	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
meloxicam	citalopram

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
meloxicam	fluoxetine
meloxicam	fluvoxamine
meloxicam	paroxetine
meloxicam	6-chloro-2-[piperidinyl-4-thio]pyridine
meloxicam	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
meloxicam	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
meloxicam	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
meloxicam	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
meloxicam	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
parecoxib	citalopram
parecoxib	fluoxetine
parecoxib	fluvoxamine
parecoxib	paroxetine
parecoxib	6-chloro-2-[piperidinyl-4-thio]pyridine
parecoxib	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
parecoxib	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
parecoxib	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
parecoxib	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
parecoxib	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
4-(4-cyclohexyl-2-methyloxazol-5-yl)-2-fluorobenzenesulfonamide	citalopram
4-(4-cyclohexyl-2-methyloxazol-5-yl)-2-fluorobenzenesulfonamide	fluoxetine
4-(4-cyclohexyl-2-methyloxazol-5-yl)-2-fluorobenzenesulfonamide	fluvoxamine
4-(4-cyclohexyl-2-methyloxazol-5-yl)-2-fluorobenzenesulfonamide	paroxetine
4-(4-cyclohexyl-2-methyloxazol-5-yl)-2-fluorobenzenesulfonamide	6-chloro-2-[piperidinyl-4-thio]pyridine
4-(4-cyclohexyl-2-methyloxazol-5-yl)-2-fluorobenzenesulfonamide	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
4-(4-cyclohexyl-2-methyloxazol-5-yl)-2-fluorobenzenesulfonamide	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
4-(4-cyclohexyl-2-methyloxazol-5-yl)-2-fluorobenzenesulfonamide	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
4-(4-cyclohexyl-2-methyloxazol-5-yl)-2-fluorobenzenesulfonamide	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
4-(4-cyclohexyl-2-methyloxazol-5-yl)-2-fluorobenzenesulfonamide	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
2-(3,5-difluorophenyl)-3-(4-(methylsulfonyl)phenyl)-2-cyclopenten-1-one	citalopram
2-(3,5-difluorophenyl)-3-(4-(methylsulfonyl)phenyl)-2-cyclopenten-1-one	fluoxetine
2-(3,5-difluorophenyl)-3-(4-(methylsulfonyl)phenyl)-2-cyclopenten-1-one	fluvoxamine
2-(3,5-difluorophenyl)-3-(4-(methylsulfonyl)phenyl)-2-cyclopenten-1-one	paroxetine
2-(3,5-difluorophenyl)-3-(4-(methylsulfonyl)phenyl)-2-cyclopenten-1-one	6-chloro-2-[piperidinyl-4-thio]pyridine
2-(3,5-difluorophenyl)-3-(4-(methylsulfonyl)phenyl)-2-cyclopenten-1-one	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
2-(3,5-difluorophenyl)-3-(4-(methylsulfonyl)phenyl)-2-cyclopenten-1-one	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
2-(3,5-difluorophenyl)-3-(4-(methylsulfonyl)phenyl)-2-cyclopenten-1-one	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
2-(3,5-difluorophenyl)-3-(4-(methylsulfonyl)phenyl)-2-cyclopenten-1-one	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
2-(3,5-difluorophenyl)-3-(4-(methylsulfonyl)phenyl)-2-cyclopenten-1-one	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
N-[2-(cyclohexyloxy)-4-nitrophenyl]methanesulfonamide	citalopram
N-[2-(cyclohexyloxy)-4-nitrophenyl]methanesulfonamide	fluoxetine
N-[2-(cyclohexyloxy)-4-nitrophenyl]methanesulfonamide	fluvoxamine
N-[2-(cyclohexyloxy)-4-nitrophenyl]methanesulfonamide	paroxetine
N-[2-(cyclohexyloxy)-4-nitrophenyl]methanesulfonamide	6-chloro-2-[piperidinyl-4-thio]pyridine
N-[2-(cyclohexyloxy)-4-nitrophenyl]methanesulfonamide	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
N-[2-(cyclohexyloxy)-4-nitrophenyl]methanesulfonamide	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
N-[2-(cyclohexyloxy)-4-nitrophenyl]methanesulfonamide	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
N-[2-(cyclohexyloxy)-4-nitrophenyl]methanesulfonamide	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
N-[2-(cyclohexyloxy)-4-nitrophenyl]methanesulfonamide	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
2-(3,4-difluorophenyl)-4-(3-hydroxy-3-methylbutoxy)-5-[4-(methylsulfonyl)phenyl]-3(2H)-pyridazinone	citalopram
2-(3,4-difluorophenyl)-4-(3-hydroxy-3-methylbutoxy)-5-[4-(methylsulfonyl)phenyl]-3(2H)-pyridazinone	fluoxetine
2-(3,4-difluorophenyl)-4-(3-hydroxy-3-methylbutoxy)-5-[4-(methylsulfonyl)phenyl]-3(2H)-pyridazinone	fluvoxamine
2-(3,4-difluorophenyl)-4-(3-hydroxy-3-methylbutoxy)-5-[4-(methylsulfonyl)phenyl]-3(2H)-pyridazinone	paroxetine
2-(3,4-difluorophenyl)-4-(3-hydroxy-3-methylbutoxy)-5-[4-(methylsulfonyl)phenyl]-3(2H)-pyridazinone	6-chloro-2-[piperidinyl-4-thio]pyridine
2-(3,4-difluorophenyl)-4-(3-hydroxy-3-methylbutoxy)-5-[4-(methylsulfonyl)phenyl]-3(2H)-pyridazinone	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
2-(3,4-difluorophenyl)-4-(3-hydroxy-3-methylbutoxy)-5-[4-(methylsulfonyl)phenyl]-3(2H)-pyridazinone	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
2-(3,4-difluorophenyl)-4-(3-hydroxy-3-methylbutoxy)-5-[4-(methylsulfonyl)phenyl]-3(2H)-pyridazinone	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
2-(3,4-difluorophenyl)-4-(3-hydroxy-3-methylbutoxy)-5-[4-(methylsulfonyl)phenyl]-3(2H)-pyridazinone	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
2-(3,4-difluorophenyl)-4-(3-hydroxy-3-methylbutoxy)-5-[4-(methylsulfonyl)phenyl]-3(2H)-pyridazinone	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
2-[(2,4-dichloro-6-methylphenyl)amino]-5-ethyl-benzeneacetic acid	citalopram
2-[(2,4-dichloro-6-methylphenyl)amino]-5-ethyl-benzeneacetic acid	fluoxetine

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
2-[(2,4-dichloro-6-methylphenyl)amino]-5-ethyl-benzeneacetic acid	fluvoxamine
2-[(2,4-dichloro-6-methylphenyl)amino]-5-ethyl-benzeneacetic acid	paroxetine
2-[(2,4-dichloro-6-methylphenyl)amino]-5-ethyl-benzeneacetic acid	6-chloro-2-[piperidinyl-4-thio]pyridine
2-[(2,4-dichloro-6-methylphenyl)amino]-5-ethyl-benzeneacetic acid	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
2-[(2,4-dichloro-6-methylphenyl)amino]-5-ethyl-benzeneacetic acid	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
2-[(2,4-dichloro-6-methylphenyl)amino]-5-ethyl-benzeneacetic acid	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
2-[(2,4-dichloro-6-methylphenyl)amino]-5-ethyl-benzeneacetic acid	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
2-[(2,4-dichloro-6-methylphenyl)amino]-5-ethyl-benzeneacetic acid	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
(3Z)-3-[(4-chlorophenyl)[4-(methylsulfonyl)phenyl]methylene]dihydro-2(3H)-furanone	citalopram
(3Z)-3-[(4-chlorophenyl)[4-(methylsulfonyl)phenyl]methylene]dihydro-2(3H)-furanone	fluoxetine
(3Z)-3-[(4-chlorophenyl)[4-(methylsulfonyl)phenyl]methylene]dihydro-2(3H)-furanone	fluvoxamine
(3Z)-3-[(4-chlorophenyl)[4-(methylsulfonyl)phenyl]methylene]dihydro-2(3H)-furanone	paroxetine
(3Z)-3-[(4-chlorophenyl)[4-(methylsulfonyl)phenyl]methylene]dihydro-2(3H)-furanone	6-chloro-2-[piperidinyl-4-thio]pyridine
(3Z)-3-[(4-chlorophenyl)[4-(methylsulfonyl)phenyl]methylene]dihydro-2(3H)-furanone	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
(3Z)-3-[(4-chlorophenyl)[4-(methylsulfonyl)phenyl]methylene]dihydro-2(3H)-furanone	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
(3Z)-3-[(4-chlorophenyl)[4-(methylsulfonyl)phenyl]methylene]dihydro-2(3H)-furanone	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
(3Z)-3-[(4-chlorophenyl)[4-(methylsulfonyl)phenyl]methylene]dihydro-2(3H)-furanone	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
(3Z)-3-[(4-chlorophenyl)[4-(methylsulfonyl)phenyl]methylene]dihydro-2(3H)-furanone	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
(S)-6,8-dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid	citalopram

Cyclooxygenase-2 Selective Inhibitor	Serotonin-Modulating Agent
(S)-6,8-dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid	fluoxetine
(S)-6,8-dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid	fluvoxamine
(S)-6,8-dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid	paroxetine
(S)-6,8-dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid	6-chloro-2-[piperidinyl-4-thio]pyridine
(S)-6,8-dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
(S)-6,8-dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
(S)-6,8-dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
(S)-6,8-dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
(S)-6,8-dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide
lumiracoxib	citalopram
lumiracoxib	fluoxetine
lumiracoxib	fluvoxamine
lumiracoxib	paroxetine
lumiracoxib	6-chloro-2-[piperidinyl-4-thio]pyridine
lumiracoxib	8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione
lumiracoxib	1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine
lumiracoxib	3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol
lumiracoxib	4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile
lumiracoxib	3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl)phenyl]benzamide

DIAGNOSIS OF A VASO-OCCLUSION

[0660] One aspect of the invention encompasses diagnosing a subject in need of treatment or prevention for a vaso-occlusive event. A number of suitable methods for diagnosing a vaso-occlusion may be used in the practice of the invention. In one such method, ultrasound may be employed. This method examines the blood flow in the major arteries and veins in the arms and legs with the use of ultrasound (high-frequency sound waves). In one embodiment, the test may combine Doppler[®] ultrasonography, which uses audio measurements to "hear" and measure the blood flow and duplex ultrasonography, which provides a visual image. In an alternative embodiment, the test may utilize multifrequency ultrasound or multifrequency transcranial Doppler[®] (MTCD) ultrasound.

[0661] Another method that may be employed encompasses injection of the subject with a compound that can be imaged. In one alternative of this embodiment, a small amount of radioactive material is injected into the subject and then standard techniques that rely on monitoring blood flow to detect a blockage, such as magnetic resonance direct thrombus imaging (MRDTI), may be utilized to image the vaso-occlusion. In an alternative embodiment, ThromboView[®] (commercially available from Agenix Limited) uses a clot-binding monoclonal antibody attached to a radiolabel. In addition to the methods identified herein, a number of other suitable methods known in the art for diagnosis of vaso-occlusive events may be utilized.

INDICATIONS TO BE TREATED

[0662] Generally speaking, the composition comprising a therapeutically effective amount of a cyclooxygenase-2 selective inhibitor and a therapeutically effective amount of a serotonin-modulating agent may be employed to treat a number of conditions resulting from a reduction in blood flow to the central nervous system.

[0663] In some aspects, the invention provides a method to treat a central nervous system cell to prevent damage in response to a decrease in blood flow to the cell. Typically the severity of damage that may be prevented will depend in large part on the degree of reduction in blood flow to the cell and the duration of the reduction. By way of example, the normal amount of perfusion to brain gray matter in humans is about 60 to 70 mL/100 g of brain tissue/min. Death of central nervous system cells typically occurs when the flow of blood falls below approximately 8-10 mL/100 g of brain tissue/min, while at slightly higher levels (i.e. 20-35 mL/100 g of brain tissue/min) the tissue remains alive but not able to function. In one embodiment, apoptotic or necrotic cell death may be prevented. In still a further embodiment, ischemic-mediated damage, such as cytotoxic edema or central nervous system tissue anoxemia, may be prevented. In each embodiment, the central nervous system cell may be a spinal cell or a brain cell.

[0664] Another aspect of this invention encompasses administering the composition to a subject to treat a central nervous system ischemic condition. Any central nervous system ischemic condition may be treated by the composition of the invention. In one embodiment, the ischemic condition is a stroke that results in any type of ischemic central nervous system damage, such as apoptotic or necrotic cell death, cytotoxic edema or central nervous system tissue anoxemia. The stroke may impact any area of the brain or be caused by any etiology commonly known to result in the occurrence of a stroke. In one alternative of this embodiment, the stroke is a brain stem stroke. Generally speaking,

brain stem strokes strike the brain stem, which control involuntary life-support functions such as breathing, blood pressure, and heartbeat. In another alternative of this embodiment, the stroke is a cerebellar stroke. Typically, cerebellar strokes impact the cerebellum area of the brain, which controls balance and coordination. In still another embodiment, the stroke is an embolic stroke. In general terms, embolic strokes may impact any region of the brain and typically result from the blockage of an artery by a vaso-occlusion. In yet another alternative, the stroke may be a hemorrhagic stroke. Like embolic strokes, hemorrhagic stroke may impact any region of the brain, and typically result from a ruptured blood vessel characterized by a hemorrhage (bleeding) within or surrounding the brain. In a further embodiment, the stroke is a thrombotic stroke. Typically, thrombotic strokes result from the blockage of a blood vessel by accumulated deposits.

[0665] In another embodiment, the ischemic condition may result from a disorder that occurs in a part of the subject's body outside of the central nervous system, but yet still causes a reduction in blood flow to the central nervous system. These disorders may include, but are not limited to a peripheral vascular disorder, a venous thrombosis, a pulmonary embolus, a myocardial infarction, a transient ischemic attack, unstable angina, or sickle cell anemia. Moreover, the central nervous system ischemic condition may occur as result of the subject undergoing a surgical procedure. By way of example, the subject may be undergoing heart surgery, lung surgery, spinal surgery, brain surgery, vascular surgery, abdominal surgery, or organ transplantation surgery. The organ transplantation surgery may include heart, lung, pancreas or liver transplantation surgery. Moreover, the central nervous system ischemic condition may occur as a result of a trauma or injury to a part of the subject's body outside the central nervous system. By way of example the trauma or injury may cause a degree of bleeding that significantly reduces the total volume of blood in the subject's body. Because of this reduced total volume, the amount of blood flow to the central nervous system is concomitantly reduced. By way of further example, the trauma or injury may also result in the formation of a vaso-occlusion that restricts blood flow to the central nervous system.

[0666] Of course it is contemplated that the composition may be employed to treat any central nervous system ischemic condition irrespective of the cause of the condition. In one embodiment, the ischemic condition results from a vaso-occlusion. The vaso-occlusion may be any type of occlusion, but is typically a cerebral thrombosis or a cerebral embolism. In a further embodiment, the ischemic condition may result from a hemorrhage. The hemorrhage may be any type of hemorrhage, but is generally a cerebral

hemorrhage or a subarachnoid hemorrhage. In still another embodiment, the ischemic condition may result from the narrowing of a vessel. Generally speaking, the vessel may narrow as a result of a vasoconstriction such as occurs during vasospasms, or due to arteriosclerosis. In yet another embodiment, the ischemic condition results from an injury to the brain or spinal cord.

[0667] In yet another aspect, the composition is administered to reduce infarct size of the ischemic core following a central nervous system ischemic condition. Moreover, the composition may also be beneficially administered to reduce the size of the ischemic penumbra or transitional zone following a central nervous system ischemic condition.

[0668] In addition to a cyclooxygenase-2 selective inhibitor and a serotonin-modulating agent, the composition of the invention may also include any agent that ameliorates the effect of a reduction in blood flow to the central nervous system. In one embodiment, the agent is an anticoagulant including thrombin inhibitors such as heparin and Factor Xa inhibitors such as warafin. In another embodiment, the agent is a thrombolytic agent including tissue plasminogen activator, urokinase, or desmoteplase (vampire bat plasminogen activator). In an additional embodiment, the agent is an anti-platelet inhibitor such as a GP IIb/IIIa inhibitor. Additional agents include but are not limited to, HMG-CoA synthase inhibitors; squalene epoxidase inhibitors; squalene synthetase inhibitors (also known as squalene synthase inhibitors); acyl-coenzyme A; cholesterol acyltransferase (ACAT) inhibitors; probucol; niacin; fibrates such as clofibrate, fenofibrate, and gemfibrozil; cholesterol absorption inhibitors; bile acid sequestrants; LDL (low density lipoprotein) receptor inducers; vitamin B₆ (also known as pyridoxine) and the pharmaceutically acceptable salts thereof such as the HCl salt; vitamin B₁₂ (also known as cyanocobalamin); β -adrenergic receptor blockers; folic acid or a pharmaceutically acceptable salt or ester thereof such as the sodium salt and the methylglucamine salt; and anti-oxidant vitamins such as vitamin C and E and beta carotene.

[0669] In a further aspect, the composition may be employed to reverse or lessen central nervous system cell damage following a traumatic brain or spinal cord injury. Traumatic brain or spinal cord injury may result from a wide variety of causes including, for example, blows to the head or back from objects; penetrating injuries from missiles, bullets, and shrapnel; falls; skull fractures with resulting penetration by bone pieces; and sudden acceleration or deceleration injuries. The composition of the invention may be beneficially utilized to treat the traumatic injury irrespective of its cause.

[0670] The composition may also beneficially be employed to increase recovery of neural cell function following brain or spinal cord injury. Generally speaking, when neurons are lost due to disease or trauma, they are not replaced. Rather, the remaining neurons must adapt to whatever loss occurred by altering their function or functional relationship relative to other neurons. Following injury, neural tissue begins to produce trophic repair factors, such as nerve growth factor and neuron cell adhesion molecules, which retard further degeneration and promote synaptic maintenance and the development of new synaptic connections. But, as the lost cells are not replaced, existing cells must take over some of the functions of the missing cells, i.e., they must "learn" to do something new. In part, recovery of function from brain traumatic damage involves plastic changes that occur in brain structures other than those damaged. Indeed, in many cases, recovery from brain damage represents the taking over by healthy brain regions of the functions of the damaged area. Thus the composition of the present invention may be administered to facilitate learning of new functions by uninjured brain areas to compensate for the loss of function by other regions.

EXAMPLES

[0671] A combination therapy of a COX-2 selective inhibitor and a serotonin-modulating agent for the treatment or prevention of a vaso-occlusive event or a related disorder in a subject can be evaluated as described in the following tests detailed below.

[0672] A particular combination therapy comprising a serotonin-modulating agent and a COX-2 inhibitor can be evaluated in comparison to a control treatment such as a placebo treatment, administration of a COX-2 inhibitor only or administration of a serotonin-modulating agent only. By way of example, a combination therapy may contain any of the serotonin-modulating agents and any of the COX-2 inhibitors detailed in the present invention, including the combinations set forth in Tables 4a, 4b, or 4c. The dosages of a serotonin-modulating agent and a COX-2 inhibitor in a particular therapeutic combination may be readily determined by a skilled artisan conducting the study. The length of the study treatment will vary on a particular study and can also be determined by one of ordinary skill in the art. By way of example, the combination therapy may be administered for 4 weeks. The serotonin-modulating agent and COX-2 inhibitor can be administered by any route as described herein, but are preferably administered orally for human subjects.

EXAMPLE 1 - EVALUATION OF COX-1 AND COX-2 ACTIVITY *IN VITRO*

[0673] The COX-2 inhibitors suitable for use in this invention exhibit selective inhibition of COX-1 over COX-2, as measured by IC_{50} values when tested *in vitro* according to the following activity assays.

PREPARATION OF RECOMBINANT COX BACULOVIRUSES

[0674] Recombinant COX-1 and COX-2 are prepared as described by Gierse et al, [*J. Biochem.*, 305, 479-84 (1995)]. A 2.0 kb fragment containing the coding region of either human or murine COX-1 or human or murine COX-2 is cloned into a BamH1 site of the baculovirus transfer vector pVL1393 (Invitrogen) to generate the baculovirus transfer vectors for COX-1 and COX-2 in a manner similar to the method of D.R. O'Reilly et al (*Baculovirus Expression Vectors: A Laboratory Manual* (1992)). Recombinant baculoviruses are isolated by transfecting 4 μ g of baculovirus transfer vector DNA into SF9 insect cells (2×10^8) along with 200 ng of linearized baculovirus plasmid DNA by the calcium phosphate method. See M.D. Summers and G.E. Smith, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Agric. Exp. Station Bull. 1555 (1987). Recombinant viruses are purified by three rounds of plaque purification and high titer (10^7 - 10^8 pfu/mL) stocks of virus are prepared. For large scale production, SF9 insect cells are infected in 10 liter fermentors (0.5×10^6 /mL) with the recombinant baculovirus stock such that the multiplicity of infection is 0.1. After 72 hours the cells are centrifuged and the cell pellet is homogenized in Tris/Sucrose (50 mM: 25%, pH 8.0) containing 1% 3-[(3-cholamidopropyl)-dimethylammonio]-1-propanesulfonate (CHAPS). The homogenate is centrifuged at 10,000xG for 30 minutes, and the resultant supernatant is stored at -80°C before being assayed for COX activity.

ASSAY FOR COX-1 AND COX-2 ACTIVITY

[0675] COX activity is assayed as PGE2 formed/ μ g protein/time using an ELISA to detect the prostaglandin released. CHAPS-solubilized insect cell membranes containing the appropriate COX enzyme are incubated in a potassium phosphate buffer (50 mM, pH 8.0) containing epinephrine, phenol, and heme with the addition of arachidonic acid (10 μ M). Compounds are pre-incubated with the enzyme for 10-20 minutes prior to the addition of arachidonic acid. Any reaction between the arachidonic acid and the enzyme is stopped after ten minutes at 37°C by transferring 40 μ l of reaction mix into 160 μ l ELISA buffer and 25 μ M indomethacin. The PGE2 formed is measured by standard ELISA technology (Cayman Chemical).

FAST ASSAY FOR COX-1 AND COX-2 ACTIVITY

[0676] COX activity is assayed as PGE₂ formed/ μ g protein/time using an ELISA to detect the prostaglandin released. CHAPS-solubilized insect cell membranes containing the appropriate COX enzyme are incubated in a potassium phosphate buffer (0.05 M Potassium phosphate, pH 7.5, 2 μ M phenol, 1 μ M heme, 300 μ M epinephrine) with the addition of 20 μ l of 100 μ M arachidonic acid (10 μ M). Compounds are pre-incubated with the enzyme for 10 minutes at 25°C prior to the addition of arachidonic acid. Any reaction between the arachidonic acid and the enzyme is stopped after two minutes at 37°C by transferring 40 μ l of reaction mix into 160 μ l ELISA buffer and 25 μ M indomethacin. Indomethacin, a non-selective COX-2/COX-1 inhibitor, may be utilized as a positive control. The PGE₂ formed is typically measured by standard ELISA technology utilizing a PGE₂ specific antibody, available from a number of commercial sources.

[0677] Each compound to be tested may be individually dissolved in 2 ml of dimethyl sulfoxide (DMSO) for bioassay testing to determine the COX-1 and COX-2 inhibitory effects of each particular compound. Potency is typically expressed by the IC₅₀ value expressed as g compound/ml solvent resulting in a 50% inhibition of PGE₂ production. Selective inhibition of COX-2 may be determined by the IC₅₀ ratio of COX-1/COX-2.

[0678] By way of example, a primary screen may be performed in order to determine particular compounds that inhibit COX-2 at a concentration of 10 μ g/ml. The compound may then be subjected to a confirmation assay to determine the extent of COX-2 inhibition at three different concentrations (e.g., 10 μ g/ml, 3.3 μ g/ml and 1.1 μ g/ml). After this screen, compounds can then be tested for their ability to inhibit COX-1 at a concentration of 10 μ g/ml. With this assay, the percentage of COX inhibition compared to control can be determined, with a higher percentage indicating a greater degree of COX inhibition. In addition, the IC₅₀ value for COX-1 and COX-2 can also be determined for the tested compound. The selectivity for each compound may then be determined by the IC₅₀ ratio of COX-1/COX-2, as set-forth above.

EXAMPLE 2 - METHODS FOR MEASURING PLATELET AGGREGATION AND PLATELET ACTIVATION MARKERS

[0679] The following studies can be performed in human subjects or laboratory animal models, such as mice. Prior to the initiation of a clinical study involving human subjects, the study should be approved by the appropriate Human Subjects Committee

and subjects should be informed about the study and give written consent prior to participation.

[0680] Platelet activation can be determined by a number of tests available in the art. Several such tests are described below. In order to determine the effectiveness of the treatment, the state of platelet activation is evaluated at several time points during the study, such as before administering the combination treatment and once a week during treatment. The exemplary procedures for blood sampling and the analyses that can be used to monitor platelet aggregation are listed below.

PLATELET AGGREGATION STUDY

[0681] Blood samples are collected from an antecubital vein via a 19-gauge needle into two plastic tubes. Each sample of free flowing blood is collected through a fresh venipuncture site distal to any intravenous catheters using a needle and Vacutainer hood into 7 cc vacutainer tubes (one with CTAD (dipyridamole), and the other with 3.8% trisodium citrate). If blood is collected simultaneously for any other studies, it is preferable that the platelet sample be obtained second or third, but not first. If only the platelet sample is collected, the initial 2-3 cc of blood is discharged and then the vacutainer tube is filled. The venipuncture is adequate if the tube fills within 15 seconds. All collections are performed by trained personnel.

[0682] After the blood samples for each subject have been collected into two Vacutainer tubes, they are immediately, but gently, inverted 3 to 5 times to ensure complete mixing of the anticoagulant. Tubes are not shaken. The Vacutainer tubes are filled to capacity, since excess anticoagulant can alter platelet function. Attention is paid to minimizing turbulence whenever possible. Small steps, such as slanting the needle in the Vacutainer to have the blood run down the side of tube instead of shooting all the way to the bottom, can result in significant improvement. These tubes are kept at room temperature and transferred directly to the laboratory personnel responsible for preparing the samples. The Vacutainer tubes are not chilled at any time.

[0683] Trisodium citrate (3.8%) and whole blood is immediately mixed in a 1:9 ratio, and then centrifuged at 1200 g for 2.5 minutes, to obtain platelet-rich plasma (PRP), which is kept at room temperature for use within 1 hour for platelet aggregation studies. Platelet count is determined in each PRP sample with a Coulter Counter ZM (Coulter Co., Hialeah, Fla.). Platelet numbers are adjusted to 3.50×10^8 /ml for aggregation with homologous platelet-poor plasma. PRP and whole blood aggregation tests are performed simultaneously. Whole blood is diluted 1:1 with the 0.5 ml PBS, and then swirled gently to

mix. The cuvette with the stirring bar is placed in the incubation well and allowed to warm to 37°C for 5 minutes. Then the samples are transferred to the assay well. An electrode is placed in the sample cuvette. Platelet aggregation is stimulated with 5 μ M ADP, 1 μ g/ml collagen, and 0.75 mM arachidonic acid. All agonists are obtained, e.g., from Chronolog Corporation (Hawertown, Pa.). Platelet aggregation studies are performed using a Chrono-Log Whole Blood Lumi-Aggregometer (model 560-Ca). Platelet aggregability is expressed as the percentage of light transmittance change from baseline using platelet-poor plasma as a reference at the end of recording time for plasma samples, or as a change in electrical impedance for whole blood samples. Aggregation curves are recorded for 4 minutes and analyzed according to internationally established standards using Aggrolink[®] software.

[0684] Aggregation curves of subjects receiving a combination therapy containing a serotonin-modulating agent and a COX-2 inhibitor can then be compared to the aggregation curves of subjects receiving a control treatment in order to determine the efficacy of said combination therapy.

WASHED PLATELETS FLOW CYTOMETRY

[0685] Venous blood (8 ml) is collected in a plastic tube containing 2 ml of acid-citrate-dextrose (ACD) (7.3 g citric acid, 22.0 g sodium citrate x 2H₂O and 24.5 glucose in 1000 ml distilled water) and mixed well. The blood-ACD mixture is centrifuged at 1000 r.p.m. for 10 minutes at room temperature. The upper 2/3 of the platelet-rich plasma (PRP) is then collected and adjusted to pH=6.5 by adding ACD. The PRP is then centrifuged at 3000 r.p.m. for 10 minutes. The supernatant is removed and the platelet pellet is gently resuspended in 4 cc of the washing buffer (10 mM Tris/HCl, 0.15 M NaCl, 20 mM EDTA, pH=7.4). Platelets are washed in the washing buffer, and in TBS (10 mM Tris, 0.15 M NaCl, pH=7.4). All cells are then divided into the appropriate number of tubes. By way of example, if 9 different surface markers are evaluated, as described herein, then the cells should be divided into ten tubes, such that nine tubes containing washed platelets are incubated with 5 μ l fluorescein isothiocyanate (FITC)-conjugated antibodies in the dark at +4°C for 30 minutes, and one tube remains unstained and serves as a negative control. Surface antigen expression is measured with monoclonal murine anti-human antibodies, such as CD9 (p24); CD41a (IIb/IIIa, aIIbb3); CD42b (Ib); CD61(IIIa) (DAKO Corporation, Carpinteria, Calif.); CD49b (VLA-2, or a2b1); CD62p (P-selectin); CD31 (PECAM-1); CD 41b (IIb); and CD51/CD61 (vitronectin receptor, avb3) (PharMingen, San Diego Calif.), as the expression of these antigens on the cells is

associated with platelet activation. After incubation, the cells are washed with TBS and resuspended in 0.25 ml of 1% paraformaldehyde. Samples are stored in the refrigerator at +4°C, and analyzed on a Becton Dickinson FACScan flow cytometer with laser output of 15 mw, excitation at 488 nm, and emission detection at 530+/-30 nm. The data can be collected and stored in list mode, and then analyzed using CELLQuest® software. FACS procedures are described in detail in, e.g., Gurbel, P. A. et al., *J Amer Coll Cardiol* 31: 1466-1473 (1998); Serebruany, V. L. et al., *Am Heart J* 136: 398-405 (1998); Gurbel, P. A. et al., *Coron Artery Dis* 9: 451-456 (1998) and Serebruany, V. L. et al., *Arterioscl Thromb Vasc Biol* 19: 153-158 (1999).

[0686] The antibody staining of platelets isolated from subjects receiving a combination therapy can then be compared to the staining of platelets isolated from subjects receiving a control treatment in order to determine the effect of the combination therapy on platelets.

WHOLE BLOOD FLOW CYTOMETRY

[0687] Four cc of blood is collected in a tube, containing 2 cc of acid-citrate-dextrose (ACD, see previous example) and mixed well. The buffer, TBS (10 mM Tris, 0.15 M NaCl, pH 7.4) and the following fluorescein isothiocyanate (FITC) conjugated monoclonal antibodies (PharMingen, San Diego, Calif., USA, and DAKO, Calif., USA) are removed from a refrigerator and allowed to warm at room temperature (RT) prior to their use. The non-limiting examples of antibodies that can be used include CD41 (IIb/IIIa), CD31 (PECAM-1), CD62p (P-selectin), and CD51/61 (Vitronectin receptor). For each subject, six amber tubes (1.25 ml) are one Eppendorf tube (1.5 ml) are obtained and marked appropriately. 450 µl of TBS buffer is pipetted to the labeled Eppendorf tube. A patient's whole blood tube is inverted gently twice to mix, and 50 µl of whole blood is pipetted to the appropriately labeled Eppendorf tube. The Eppendorf tube is capped and the diluted whole blood is mixed by inverting the Eppendorf tube gently two times, followed by pipetting 50 µl of diluted whole blood to each amber tube. 5 µl of appropriate antibody is pipetted to the bottom of the corresponding amber tube. The tubes are covered with aluminum foil and incubated at 4°C for 30 minutes. After incubation, 400 µl of 2% buffered paraformaldehyde is added. The amber tubes are closed with a lid tightly and stored in a refrigerator at 4°C until the flow cytometric analysis. The samples are analyzed on a Becton Dickinson FACScan flow cytometer. These data are collected in list mode files and then analyzed. As mentioned in (B.), the antibody staining of platelets isolated from

subjects receiving a combination therapy can then be compared to the staining of platelets isolated from subjects receiving a control treatment.

ELISA

[0688] Enzyme-linked immunosorbent assays (ELISA) are used according to standard techniques and as described herein. Eicosanoid metabolites may be used to determine platelet aggregation. The metabolites are analyzed due to the fact that eicosanoids have a short half-life under physiological conditions. Thromboxane B₂ (TXB₂), the stable breakdown product of thromboxane A₂ and 6keto-PGF₁ alpha, the stable degradation product of prostacyclin may be tested. Thromboxane B₂ is a stable hydrolysis product of TXA₂ and is produced following platelet aggregation induced by a variety of agents, such as thrombin and collagen. 6keto-prostaglandin F₁ alpha is a stable hydrolyzed product of unstable PGI₂ (prostacyclin). Prostacyclin inhibits platelet aggregation and induces vasodilation. Thus, quantitation of prostacyclin production can be made by determining the level of 6keto-PGF₁. The metabolites may be measured in the platelet poor plasma (PPP), which is kept at -4°C. Also, plasma samples may also be extracted with ethanol and then stored at -80° C before final prostaglandin determination, using, e.g., TiterZymes[®] enzyme immunoassays according to standard techniques (PerSeptive Diagnostics, Inc., Cambridge, Mass., USA). ELISA kits for measuring TXB₂ and 6keto-PGF₁ are also commercially available.

[0689] The amounts of TXB₂ and 6keto-PGF₁ in plasma of subjects receiving a combination therapy and subjects receiving a control therapy can be compared to determine the efficacy of the combination treatment.

CLOSURE TIME MEASURED WITH THE DADE BEHRING PLATELET FUNCTION ANALYZER, PFA-100[®]

[0690] PFA-100[®] can be used as an *in vitro* system for the detection of platelet dysfunction. It provides a quantitative measure of platelet function in anticoagulated whole blood. The system comprises a microprocessor-controlled instrument and a disposable test cartridge containing a biologically active membrane. The instrument aspirates a blood sample under constant vacuum from the sample reservoir through a capillary and a microscopic aperture cut into the membrane. The membrane is coated with collagen and epinephrine or adenosine 5'-diphosphate. The presence of these biochemical stimuli, and the high shear rates generated under the standardized flow conditions, result in platelet attachment, activation, and aggregation, slowly building a stable platelet plug at the

aperture. The time required to obtain full occlusion of the aperture is reported as the "closure time," which normally ranges from one to three minutes.

[0691] The membrane in the PFA-100[®] test cartridge serves as a support matrix for the biological components and allows placement of the aperture. The membrane is a standard nitrocellulose filtration membrane with an average pore size of 0.45 μm . The blood entry side of the membrane was coated with 2 μg of fibrillar Type I equine tendon collagen and 10 μg of epinephrine bitartrate or 50 μg of adenosine 5'-diphosphate (ADP). These agents provide controlled stimulation to the platelets as the blood sample passes through the aperture. The collagen surface also served as a well-defined matrix for platelet deposition and attachment.

[0692] The principle of the PFA-100[®] test is very similar to that described by Kratzer and Born (Kratzer, et al., *Haemostasis* 15: 357-362 (1985)). The test utilizes whole blood samples collected in 3.8% of 3.2% sodium citrate anticoagulant. The blood sample is aspirated through the capillary into the cup where it comes in contact with the coated membrane, and then passes through the aperture. In response to the stimulation by collagen and epinephrine or ADP present in the coating, and the shear stresses at the aperture, platelets adhere and aggregate on the collagen surface starting at the area surrounding the aperture. During the course of the measurement, a stable platelet plug forms that ultimately occludes the aperture. The time required to obtain full occlusion of the aperture is defined as the "closure time" and is indicative of the platelet function in the sample. Accordingly, "closure times" can be compared between subjects receiving a combination therapy and the ones receiving a control therapy in order to evaluate the efficacy of the combination treatment. By way of example, a combination therapy may contain fluoxetine and celecoxib, paroxetine and valdecoxib, citalopram and rofecoxib, or escitalopram and celecoxib. It should be noted that these are only several examples, and that any of the serotonin-modulating agents and COX-2 inhibitors of the present invention may be tested as a combination therapy. The dosages of the serotonin-modulating agent and COX-2 inhibitor in a particular therapeutic combination may be readily determined by a skilled artisan conducting the study. The length of the study treatment will vary on a particular study and can also be determined by one of ordinary skill in the art. By way of example, the combination therapy may be administered for 12 weeks. The serotonin-modulating agent and COX-2 inhibitor can be administered by any route as described herein, but are preferably administered orally or intravenously for human subjects.

EXAMPLE 3 – GLOBAL ISCHEMIA AND FOCAL ISCHEMIA STUDIES

[0693] The laboratory animal study can generally be performed as described in Tanaka *et al.*, *Neurochemical Research*, Vol. 20, No. 6, 1995, pp. 663-667.

[0694] Briefly, the study can be performed with about 30 gerbils, with body weights of 65 to 80 grams. The animals are anesthetized with ketamine (100mg/kg body weight, i.p.), and silk threads are placed around both common carotid arteries without interrupting carotid artery blood flow. On the next day, bilateral common carotid arteries are exposed and then occluded with surgical clips after light ether anesthesia (see, e.g., Ogawa *et al.*, *Adv. Exp. Med. Biol.*, 287:343-347, and Ogawa *et al.*, *Brain Res.*, 591:171-175). Carotid artery blood flow is restored by releasing the clips after 5 minutes of occlusion. Body temperature is maintained about 37°C using a heating pad and an incandescent lamp. Control animals are operated on in a similar manner but the carotid arteries are not occluded. The combination therapy is administered immediately and 6 and 12 hours after recirculation in the ischemia group, whereas sham-operated animals receive placebo, which may be, e.g., the vehicle used to administer the combination therapy. Gerbils are sacrificed by decapitation 14 days after recirculation. The brain is removed rapidly and placed on crushed dry-ice to freeze the tissue.

[0695] The brain tissue can then be examined histologically for the effects of combination therapy in comparison to the placebo. For example, each brain is cut into 14 μm thick sections at -15°C. Coronal sections that include the cerebral cortex and hippocampal formation are thawed, mounted onto gelatin-coated slides, dried completely, and fixed with 10% formalin for 2 hours. The sections are stained with hematoxylin-eosin and antibodies to glial fibrillary acidic protein (GFAP), which can be commercially obtained from, e.g., Nichirei, Tokyo, Japan. Immune complexes are detected by the avidin-biotin interaction and visualized with 3,3'-diaminobenzidine tetrahydrochloride. Sections that are used as controls are stained in a similar manner without adding anti-GFAP antibody. The densities of living pyramidal cells and GFAP-positive astrocytes in the typical CA1 subfield of the hippocampus are calculated by counting the cells and measuring the total length of the CA1 cell layer in each section from 250x photomicrographs. The average densities of pyramidal cells and GFAP-positive astrocytes in the CA1 subfield for each gerbil are obtained from counting cells in one unit area in each of these sections of both left and right hemispheres.

[0696] The effects of the combination therapy in comparison with the placebo can be determined both qualitatively and quantitatively. For example, the appearance of CA1 pyramidal neurons and pyramidal cell density in the CA1 subfield may be used to

assess the efficacy of the treatment. In addition, immunohistological analysis can reveal the efficacy of combination by evaluating the presence or absence of hypertrophic GFAP-positive astrocytes in the CA1 region of treated gerbils, since the sham-operated animals should have few GFAP-positive astrocytes.

EXAMPLE 4

[0697] Rat middle cerebral artery occlusion (MCAO) models are well known in the art and useful in assessing a neuroprotective drug efficacy in stroke. By way of example, the methods and materials for MCAO model described in Turski *et al.* (*Proc. Natl. Acad. Sci. USA*, Vol. 95, pp.10960-10965, Sept. 1998) may be modified for testing the combination therapy as described above for cerebral ischemia treatment.

[0698] The permanent middle cerebral artery occlusion can be established by means of microbipolar permanent coagulation in, e.g., Fisher 344 rats (260-290 grams) anesthetized with halothane as described previously in, e.g., Lippert *et al.*, *Eur. J. Pharmacol.*, 253, pp.207-213, 1994. To determine the efficacy of the combination treatment and the therapeutic window for such treatment, the combination therapy can be administered, e.g., intravenously over 6 hours beginning 1, 2, 4, 5, 6, 7, 12, or 24 hours after MCAO. It should be noted that different doses, routes of administrations, and times of administration can also be readily tested. Furthermore, the experiment should be controlled appropriately, e.g. by administering placebo to a set of MCAO-induced rats. To evaluate the efficacy of the combination therapy, the size of infarct in the brain can be estimated stereologically, e.g., seven days after MCAO, by means of advanced image analysis.

[0699] In addition, the assessment of neuroprotective action against focal cerebral reperfusion ischemia can be performed in Wistar rats (250-300 grams) that are anesthetized with halothane and subjected to temporary occlusion of the common carotid arteries and the right middle cerebral artery (CCA/MCAO) for 90 minutes. CCAs can be occluded by means of silastic threads placed around the vessels, and MCA can be occluded by means of a steel hook attached to a micromanipulator. Blood flow stop can be verified by microscopic examination of the MCA or laser doppler flowmetry. Different doses of combination therapy can then be administered over, e.g., 6 hours starting immediately after the beginning of reperfusion or, e.g., 2 hours after the onset of reperfusion. As mentioned previously, the size of infarct in the brain can be estimated, for example, stereologically seven days after CCA/MCAO by means of image analysis.

EXAMPLE 5 – FOCAL CEREBRAL ISCHEMIA STUDY

[0700] The following procedures can be performed as described in, e.g., Nogawa *et al.*, *Journal of Neuroscience*, 17(8):2746-2755, April 15, 1997.

[0701] The middle cerebral artery (MCA) is transiently occluded in a number of Sprague Dawley rats, weighing 275-310 grams, using an intravascular occlusion model, as described in, e.g., Longa *et al.*, *Stroke* 20:84-91, 1989, Iadecola *et al.*, *Stroke* 27:1373-1380, 1996, and Zhang *et al.*, *Stroke* 27:317-323. A skilled artisan can readily determine the appropriate number of animals to be used for a particular experiment. Under halothane anesthesia (induction 5%, maintenance 1%), a 4-0 nylon monofilament with a rounded tip is inserted centripetally into the external carotid artery and advanced into the internal carotid artery until it reaches the circle of Willis. Throughout the procedure, body temperature is maintained at $37^{\circ} \pm 0.5^{\circ}\text{C}$ by a thermostatically controlled lamp. Two hours after induction of ischemia, rats are reanesthetized, and the filament is withdrawn, as described in, e.g., Zhang *et al.*, *Stroke* 27:317-323. Animals are then returned to their cages and closely monitored until recovery from anesthesia.

[0702] Under halothane anesthesia, the femoral artery is cannulated, and rats are placed on a stereotaxic frame. The arterial catheter is used for monitoring of arterial pressure and other parameters at different times after MCA occlusion. The MCA is occluded for 2 hours, as described above, and treatments are started, e.g., 6 hours after induction of ischemia. In one group of rats (e.g., 6), the combination therapy is administered, e.g., intraperitoneally, twice a day for 3 days. It should be noted that different doses, routes of administration, and times of administration can also be readily tested. A second group of rats is treated with a placebo administered in the same manner. Arterial pressure, rectal temperature, and plasma glucose are measured three times a day during the experiment. Arterial hematocrit and blood gases are measured before injection and 24, 48, and 72 hours after ischemia. Three days after MCA occlusion, brains are removed and frozen in cooled isopentane (-30°C). Coronal forebrain sections (30 μm thick) are serially cut in cryostat, collected at 300 μm intervals, and stained with thionin for determination of infarct volume by an image analyzer (e.g., MCID, Imaging Research), as described in Iadecola *et al.*, *J Cereb Blood Flow Metab*, 15:378-384, 1995. Infarct volume in cerebral cortex is corrected for swelling according to the method of Lin *et al.*, *Stroke* 24:117-121, 1993, which is based on comparing the volumes of neocortex ipsilateral and contralateral to the stroke. The correction for swelling is needed to factor out the contribution of ischemic swelling to the total volume of the lesion (see Zhang and Iadecola, *J Cereb Blood Flow Metab*, 14:574-580, 1994). Reduction of infarct size in combination

therapy-treated animals compared to animals receiving placebo is indicative of the efficacy of the combination therapy.

It should be noted that all of the above-mentioned procedures could be modified for a particular study, depending on factors such as a drug combination used, length of the study, subjects that are selected, etc. Such modifications can be designed by a skilled artisan without undue experimentation.

WHAT IS CLAIMED IS:

1. A method for treating a stroke, the method comprising:
(a) diagnosing a subject in need of treatment for a stroke; and
(b) administering to the subject a combination comprising a
cyclooxygenase-2 selective inhibitor or an isomer, a pharmaceutically acceptable salt,
5 ester, or prodrug thereof and a serotonin-modulating agent or an isomer, a
pharmaceutically acceptable salt, ester, or prodrug thereof.

2. The method of claim 1 wherein the cyclooxygenase-2 selective inhibitor has
a selectivity ratio of COX-1 IC₅₀ to COX-2 IC₅₀ not less than about 50.

3. The method of claim 1 wherein the cyclooxygenase-2 selective inhibitor has
a selectivity ratio of COX-1 IC₅₀ to COX-2 IC₅₀ not less than about 100.

4. The method of claim 1 wherein the cyclooxygenase-2 selective inhibitor is
selected from the group consisting of celecoxib, cimicoxib, deracoxib, valdecoxib,
rofecoxib, lumiracoxib, etoricoxib, meloxicam, parecoxib, 4-(4-cyclohexyl-2-methyloxazol-
5-yl)-2-fluorobenzenesulfonamide, 2-(3,5-difluorophenyl)-3-(4-(methylsulfonyl)phenyl)-2-
5 cyclopenten-1-one, N-[2-(cyclohexyloxy)-4-nitrophenyl]methanesulfonamide, 2-(3,4-
difluorophenyl)-4-(3-hydroxy-3-methylbutoxy)-5-[4-(methylsulfonyl)phenyl]-3(2H)-
pyridazinone, 2-[(2,4-dichloro-6-methylphenyl)amino]-5-ethyl-benzeneacetic acid, (3Z)-3-
[(4-chlorophenyl)[4-(methylsulfonyl)phenyl]methylene]dihydro-2(3H)-furanone, and (S)-6,8-
10 dichloro-2-(trifluoromethyl)-2H-1-benzopyran-3-carboxylic acid, or an isomer, a
pharmaceutically acceptable salt, ester, or prodrug thereof.

5. The method of claim 1 wherein the serotonin-modulating agent is selected
from the group consisting of:

citalopram;

fluoxetine;

5 fluvoxamine;

paroxetine;

norfluoxetine;

N-demethylsertraline;

escitalopram oxalate;

10 sertraline;

3aS-2-[(S)-1-azabicyclo [2.2.2] oct-3-yl]-2,3,3a,4,5,6-hexahydro-1-oxo-1H-
benz[de]isoquinoline hydrochloride;

- 3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol;
4-[3-[*t*-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile;
15 3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl) phenyl]benzamide;
N-[4-methoxy-3-(4-methyl-1-piperazinyl)phenyl]-2'-methyl-4'-(5-methyl-1,2,4-
oxadiazol-3-yl)-1,1'-biphenyl-4-carboxamide hydrochloride;
1-[(1-methylethyl)amino]-3-[2-(1H-pyrrol-1-yl)phenoxy]propan-2-ol;
1-(2-methoxyphenyl)-4-(4-succinimidobutyl)piperazine;
20 1-(2-methoxyphenyl)-4-(4-phthalimidobutyl)piperazine;
1-(1H-indol-4-yloxy)-3-[(1-methylethyl)amino]-2-propanol;
(S)-1-(1H-indol-4-yloxy)-3-[(1-methylethyl)amino]-2-propanol;
N-[3-[3-(dimethylamino)ethoxy]-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-
oxadiazol-3-yl)-[1,1'-biphenyl]-4-carboxamide;
25 8-[(2,3-dihydro-1,4-benzodioxin-2-yl)methyl]-1-phenyl-1,3,8-triazaspiro [4,5]decan-
4-one;
(S)-N-tert-butyl-3-(4-(2-methoxyphenyl)-piperazin-1-yl)-2-phenylpropanamide;
8-[3-(4-fluorophenoxy)propyl]-1-phenyl-1,3,8-triazaspiro[4.5]-decan-4-one;
N-[2-[[3-(dimethylamino)propyl]thio]phenyl]-3-phenyl-2-propenamide;
30 4-(5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-methylpiperidine;
8-chloro-11-(1-piperazinyl)-5H-dibenzo[b,e][1,4]diazepine;
4-(4-fluorobenzoyl)-1-(4-phenylbutyl)-piperidine oxalate;
3-[2-[4-(4-fluorobenzoyl)-1-piperidinyl]ethyl]-2,4[1H,3H]-quinazolinedione;
 α -phenyl-1-(2-phenylethyl)-4-piperidinemethanol;
35 metergoline phenylmethyl ester;
1,2,3,4,10,14b-hexahydro-2-methyldibenzo[c,f]pyrazino[1,2-a]azepine;
8-[5-(2,4-dimethoxy-5-(4-trifluoromethylphenylsulphonamido) phenyl-5-oxopentyl]-
1,3,8-triazaspiro[4.5]decane-2,4-dione;
N-(1-methyl-1H-indol-5-yl)-N'-3-pyridinyl-urea;
40 N-(1-methyl-5-indolyl)-N'-(3-methyl-5-isothiazolyl)urea;
(+)-cis-4,5,7a,8,9,10,11,11a-octahydro-7H-10-methylindolo[1,7-bc][2,6]-
naphthyridine;
8-[4-(4-fluorophenyl)-4-oxobutyl]-1-phenyl-1,3,8-triazaspiro[4,5]decan-4-one;
6-chloro-2-[piperidinyl-4-thio]pyridine;
45 8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione;
1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine;
5-hydroxy-3-(1-methylpiperidin-4-yl)-1H-indole;

8-[4-[4-(2-pyrimidinyl)-1-piperazinyl]butyl]-8-azaspiro[4.5]decane-7,8-dione;

5-carboxamidotryptamine maleate;

50 7-trifluoromethyl-4-(4-methyl-1-piperazinyl)pyrrolo[1,2-a]-quinoxaline;

1,4-dihydro-3-(1,2,3,6-tetrahydro-4-pyridinyl)-5H-pyrrolo[3,2-b]pyridin-5-one;

5-propoxy-3-(1,2,3,6-tetrahydro-4-pyridinyl)-1H-pyrrolo[3,2b]pyridine hydrochloride;

3-[3-(2-dimethylaminoethyl)-1H-indol-5-yl]-N-(4-methoxybenzyl)acrylamide;

8-hydroxy-2-dipropylaminotetralin hydrobromide;

55 (2R)-(+)-8-hydroxy-2-(di-n-propylamino)tetralin;

(RS)-trans-8-hydroxy-2-[N-n-propyl-N-(3'-iodo-2'-propenyl)amino]tetralin;

2-[5-[3-(4-methylsulphonylamino)benzyl-1,2,4-oxadiazol-5-yl]-1H-indol-3-

yl]ethanamine;

8-[2-(1,4-benzodioxan-2-ylmethylamino)ethyl]-8-azaspiro[4.5]decane-7,9-dione;

60 nonyloxytryptamineoxalate;

5-methoxy-3-(1,2,5,6-tetrahydro-4-pyridinyl)-1H-indole;

N-(3-trifluoromethylphenyl)piperazine;

α -methyl-5-(2-thienylmethoxy)-1H-indole-3-ethanamine;

1-(3-chlorophenyl)piperazine;


65 α -methyl-5-hydroxytryptamine maleate; and

6-chloro-2-(1-piperazinyl)pyrazine,

or is an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof.

6. A composition comprising:

(a) a cyclooxygenase-2 selective inhibitor or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof having the formula:



5 wherein:

n is an integer which is 0, 1, 2, 3 or 4;

G is O, S or NR^a;

R^a is alkyl;

R¹ is selected from the group consisting of H and aryl;

10 R² is selected from the group consisting of carboxyl, aminocarbonyl, alkylsulfonylaminocarbonyl and alkoxy carbonyl;

R^3 is selected from the group consisting of haloalkyl, alkyl, aralkyl, cycloalkyl and aryl optionally substituted with one or more radicals selected from alkylthio, nitro and alkylsulfonyl; and

- 15 each R^4 is independently selected from the group consisting of H, halo, alkyl, aralkyl, alkoxy, aryloxy, heteroaryloxy, aralkyloxy, heteroaralkyloxy, haloalkyl, haloalkoxy, alkylamino, arylamino, aralkylamino, heteroaryl amino, heteroarylalkyl amino, nitro, amino, aminosulfonyl, alkylaminosulfonyl, arylaminosulfonyl, heteroarylaminosulfonyl, aralkylaminosulfonyl, heteroaralkylaminosulfonyl, heterocyclosulfonyl, alkylsulfonyl, 20 hydroxyarylcarbonyl, nitroaryl, optionally substituted aryl, optionally substituted heteroaryl, aralkylcarbonyl, heteroarylcarbonyl, arylcarbonyl, aminocarbonyl, and alkylcarbonyl; or R^4 together with the carbon atoms to which it is attached and the remainder of ring E forms a naphthyl radical; and

(b) a serotonin-modulating agent selected from the group consisting of:

- 25 citalopram;
fluoxetine;
fluvoxamine;
paroxetine;
norfluoxetine;
30 N-demethylsertraline;
escitalopram oxalate;
sertraline;
3aS-2-[(S)-1-azabicyclo [2.2.2] oct-3-yl]-2,3,3a,4,5,6-hexahydro-1-oxo-1H-benz[de]isoquinoline hydrochloride;
35 3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol;
4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile;
3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl) phenyl]benzamide;
N-[4-methoxy-3-(4-methyl-1-piperazinyl)phenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-1,1'-biphenyl-4-carboxamide hydrochloride;
40 1-[(1-methylethyl)amino]-3-[2-(1H-pyrrol-1-yl)phenoxy]propan-2-ol;
1-(2-methoxyphenyl)-4-(4-succinimidobutyl)piperazine;
1-(2-methoxyphenyl)-4-(4-phthalimidobutyl)piperazine;
1-(1H-indol-4-yloxy)-3-[(1-methylethyl)amino]-2-propanol;
(S)-1-(1H-indol-4-yloxy)-3-[(1-methylethyl)amino]-2-propanol;
45 N-[3-[3-(dimethylamino)ethoxy]-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-[1,1'-biphenyl]-4-carboxamide;

8-[(2,3-dihydro-1,4-benzodioxin-2-yl)methyl]-1-phenyl-1,3,8-triazaspiro [4,5]decan-4-one;

(S)-N-tert-butyl-3-(4-(2-methoxyphenyl)-piperazin-1-yl)-2-phenylpropanamide;

50 8-[3-(4-fluorophenoxy)propyl]-1-phenyl-1,3,8-triazaspiro[4.5]-decan-4-one;

N-[2-[[3-(dimethylamino)propyl]thio]phenyl]-3-phenyl-2-propenamide;

4-(5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-methylpiperidine;

8-chloro-11-(1-piperazinyl)-5H-dibenzo[b,e][1,4]diazepine;

4-(4-fluorobenzoyl)-1-(4-phenylbutyl)-piperidine oxalate;

55 3-[2-[4-(4-fluorobenzoyl)-1-piperidinyl]ethyl]-2,4[1H,3H]-quinazolinedione;

α -phenyl-1-(2-phenylethyl)-4-piperidinemethanol;

metergoline phenylmethyl ester;

1,2,3,4,10,14b-hexahydro-2-methyldibenzo[c,f]pyrazino[1,2-a]azepine;

8-[5-(2,4-dimethoxy-5-(4-trifluoromethylphenylsulphonamido) phenyl-5-oxopentyl]-

60 1,3,8-triazaspiro[4.5]decane-2,4-dione;

N-(1-methyl-1H-indol-5-yl)-N'-3-pyridinyl-urea;

N-(1-methyl-5-indolyl)-N'-(3-methyl-5-isothiazolyl)urea;

(+)-cis-4,5,7a,8,9,10,11,11a-octahydro-7H-10-methylindolo[1,7-bc][2,6]-

naphthyridine;

65 8-[4-(4-fluorophenyl)-4-oxobutyl]-1-phenyl-1,3,8-triazaspiro[4,5]decan-4-one;

6-chloro-2-[piperidinyl-4-thio]pyridine;

8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione;

1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine;

5-hydroxy-3-(1-methylpiperidin-4-yl)-1H-indole;

70 8-[4-[4-(2-pyrimidinyl)-1-piperazinyl]butyl]-8-azaspiro[4.5]decane-7,8-dione;

5-carboxamidotryptamine maleate;

7-trifluoromethyl-4-(4-methyl-1-piperazinyl)pyrrolo[1,2-a]-quinoxaline;

1,4-dihydro-3-(1,2,3,6-tetrahydro-4-pyridinyl)-5H-pyrrolo[3,2-b]pyridin-5-one;

5-propoxy-3-(1,2,3,6-tetrahydro-4-pyridinyl)-1H-pyrrolo[3,2b]pyridine hydrochloride;

75 3-[3-(2-dimethylaminoethyl)-1H-indol-5-yl]-N-(4-methoxybenzyl)acrylamide;

8-hydroxy-2-dipropylaminotetralin hydrobromide;

(2R)-(+)-8-hydroxy-2-(di-n-propylamino)tetralin;

(RS)-trans-8-hydroxy-2-[N-n-propyl-N-(3'-iodo-2'-propenyl)amino]tetralin;

2-[5-[3-(4-methylsulphonylamino)benzyl-1,2,4-oxadiazol-5-yl]-1H-indol-3-


80 yl]ethanamine;

8-[2-(1,4-benzodioxan-2-ylmethylamino)ethyl]-8-azaspiro[4.5]decane-7,9-dione;

nonyloxytryptamineoxalate;
 5-methoxy-3-(1,2,5,6-tetrahydro-4-pyridinyl)-1H-indole;
 N-(3-trifluoromethylphenyl)piperazine;
 85 α -methyl-5-(2-thienylmethoxy)-1H-indole-3-ethanamine;
 1-(3-chlorophenyl)piperazine;
 α -methyl-5-hydroxytryptamine maleate; and
 6-chloro-2-(1-piperazinyl)pyrazine,
 or is an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof.

7. A composition comprising:

(a) a cyclooxygenase-2 selective inhibitor or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof of the formula:



5 wherein:

A is selected from the group consisting of a partially unsaturated or unsaturated heterocyclyl ring and a partially unsaturated or unsaturated carbocyclic ring;

R^1 is selected from the group consisting of heterocyclyl, cycloalkyl, cycloalkenyl and aryl, wherein R^1 is optionally substituted at a substitutable position with one or more
 10 radicals selected from alkyl, haloalkyl, cyano, carboxyl, alkoxycarbonyl, hydroxyl, hydroxyalkyl, haloalkoxy, amino, alkylamino, arylamino, nitro, alkoxyalkyl, alkylsulfanyl, halo, alkoxy and alkylthio;

R^2 is selected from the group consisting of methyl and amino; and

R^3 is selected from the group consisting of H, halo, alkyl, alkenyl, alkynyl, oxo,
 15 cyano, carboxyl, cyanoalkyl, heterocycloxy, alkoxy, alkylthio, alkylcarbonyl, cycloalkyl, aryl, haloalkyl, heterocyclyl, cycloalkenyl, aralkyl, heterocyclalkyl, acyl, alkylthioalkyl, hydroxyalkyl, alkoxycarbonyl, arylcarbonyl, aralkylcarbonyl, aralkenyl, alkoxyalkyl, arylthioalkyl, aryloxyalkyl, aralkylthioalkyl, aralkoxyalkyl, alkoxyaralkoxyalkyl, alkoxycarbonylalkyl, aminocarbonyl, aminocarbonylalkyl, alkylaminocarbonyl, N-
 20 arylaminocarbonyl, N-alkyl-N-arylaminocarbonyl, alkylaminocarbonylalkyl, carboxyalkyl, alkylamino, N-arylmino, N-aralkylamino, N-alkyl-N-aralkylamino, N-alkyl-N-arylmino, aminoalkyl, alkylaminoalkyl, N-arylminoalkyl, N-aralkylaminoalkyl, N-alkyl-N-aralkylaminoalkyl, N-alkyl-N-arylminoalkyl, aryloxy, aralkoxy, arylthio, aralkylthio,

alkylsulfinyl, alkylsulfonyl, aminosulfonyl, alkylaminosulfonyl, N-arylamino sulfonyl,
 25 arylsulfonyl, and N-alkyl-N-arylamino sulfonyl; and

(b) a serotonin-modulating agent selected from the group consisting of:

citalopram;

fluoxetine;

fluvoxamine;

30 paroxetine;

norfluoxetine;

N-demethylsertraline;

escitalopram oxalate;

sertraline;

35 3aS-2-[(S)-1-azabicyclo [2.2.2] oct-3-yl]-2,3,3a,4,5,6-hexahydro-1-oxo-1H-benz[de]isoquinoline hydrochloride;

3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol;

4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile;

3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl) phenyl]benzamide;

40 N-[4-methoxy-3-(4-methyl-1-piperazinyl)phenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-1,1'-biphenyl-4-carboxamide hydrochloride;

1-[(1-methylethyl)amino]-3-[2-(1H-pyrrol-1-yl)phenoxy]propan-2-ol;

1-(2-methoxyphenyl)-4-(4-succinimidobutyl)piperazine;

1-(2-methoxyphenyl)-4-(4-phthalimidobutyl)piperazine;

45 1-(1H-indol-4-yloxy)-3-[(1-methylethyl)amino]-2-propanol;

(S)-1-(1H-indol-4-yloxy)-3-[(1-methylethyl)amino]-2-propanol;

N-[3-[3-(dimethylamino)ethoxy]-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-[1,1'-biphenyl]-4-carboxamide;

8-[(2,3-dihydro-1,4-benzodioxin-2-yl)methyl]-1-phenyl-1,3,8-triazaspiro [4,5]decan-
 50 4-one;

(S)-N-tert-butyl-3-(4-(2-methoxyphenyl)-piperazin-1-yl)-2-phenylpropanamide;

8-[3-(4-fluorophenoxy)propyl]-1-phenyl-1,3,8-triazaspiro[4.5]-decan-4-one;

N-[2-[[3-(dimethylamino)propyl]thio]phenyl]-3-phenyl-2-propenamide;

4-(5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-methylpiperidine;

55 8-chloro-11-(1-piperazinyl)-5H-dibenzo[b,e][1,4]diazepine;

4-(4-fluorobenzoyl)-1-(4-phenylbutyl)-piperidine oxalate;

3-[2-[4-(4-fluorobenzoyl)-1-piperidinyl]ethyl]-2,4[1H,3H]-quinazolin-2-one;


α -phenyl-1-(2-phenylethyl)-4-piperidinemethanol;

metergoline phenylmethyl ester;

- 60 1,2,3,4,10,14b-hexahydro-2-methyldibenzo[c,f]pyrazino[1,2-a]azepine;
8-[5-(2,4-dimethoxy-5-(4-trifluoromethylphenylsulphonamido) phenyl-5-oxopentyl]-
1,3,8-triazaspiro[4.5]decane-2,4-dione;
N-(1-methyl-1H-indol-5-yl)-N'-3-pyridinyl-urea;
N-(1-methyl-5-indolyl)-N'-(3-methyl-5-isothiazolyl)urea;
- 65 (+)-cis-4,5,7a,8,9,10,11,11a-octahydro-7H-10-methylindolo[1,7-bc][2,6]-
naphthyridine;
8-[4-(4-fluorophenyl)-4-oxobutyl]-1-phenyl-1,3,8-triazaspiro[4,5]decan-4-one;
6-chloro-2-[piperidinyl-4-thio]pyridine;
8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione;
- 70 1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine;
5-hydroxy-3-(1-methylpiperidin-4-yl)-1H-indole;
8-[4-[4-(2-pyrimidinyl)-1-piperazinyl]butyl]-8-azaspiro[4.5]decane-7,8-dione;
5-carboxamidotryptamine maleate;
7-trifluoromethyl-4-(4-methyl-1-piperazinyl)pyrrolo[1,2-a]-quinoxaline;
- 75 1,4-dihydro-3-(1,2,3,6-tetrahydro-4-pyridinyl)-5H-pyrrolo[3,2-b]pyridin-5-one;
5-propoxy-3-(1,2,3,6-tetrahydro-4-pyridinyl)-1H-pyrrolo[3,2b]pyridine hydrochloride;
3-[3-(2-dimethylaminoethyl)-1H-indol-5-yl]-N-(4-methoxybenzyl)acrylamide;
8-hydroxy-2-dipropylaminotetralin hydrobromide;
(2R)-(+)-8-hydroxy-2-(di-n-propylamino)tetralin;
- 80 (RS)-trans-8-hydroxy-2-[N-n-propyl-N-(3'-iodo-2'-propenyl)amino]tetralin;
2-[5-[3-(-4-methylsulphonylamino)benzyl-1,2,4-oxadiazol-5-yl]-1H-indol-3-
yl]ethanamine;
8-[2-(1,4-benzodioxan-2-ylmethylamino)ethyl]-8-azaspiro[4.5]decane-7,9-dione;
nonyloxytryptamineoxalate;
- 85 5-methoxy-3-(1,2,5,6-tetrahydro-4-pyridinyl)-1H-indole;
N-(3-trifluoromethylphenyl)piperazine;
 α -methyl-5-(2-thienylmethoxy)-1H-indole-3-ethanamine;
1-(3-chlorophenyl)piperazine;
 α -methyl-5-hydroxytryptamine maleate; and
- 90 6-chloro-2-(1-piperazinyl)pyrazine,
or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof.

8. A composition comprising:

(a) a cyclooxygenase-2 selective inhibitor or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof having the formula:



wherein:

R¹⁶ is methyl or ethyl;

R¹⁷ is chloro or fluoro;

R¹⁸ is hydrogen or fluoro;

R¹⁹ is hydrogen, fluoro, chloro, methyl, ethyl, methoxy, ethoxy or hydroxy;

R²⁰ is hydrogen or fluoro; and

R²¹ is chloro, fluoro, trifluoromethyl or methyl, provided, however, that each of R¹⁷, R¹⁸, R²⁰ and R²¹ is not fluoro when R¹⁶ is ethyl and R¹⁹ is H; and

(b) a serotonin-modulating agent selected from the group consisting of:

citalopram;

fluoxetine;

fluvoxamine;

paroxetine;

norfluoxetine;

N-demethylsertraline;

escitalopram oxalate;

sertraline;

3aS-2-[(S)-1-azabicyclo [2.2.2] oct-3-yl]-2,3,3a,4,5,6-hexahydro-1-oxo-1H-benz[de]isoquinoline hydrochloride;

3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol;

4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile;

3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl) phenyl]benzamide;

- N-[4-methoxy-3-(4-methyl-1-piperazinyl)phenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-1,1'-biphenyl-4-carboxamide hydrochloride;
- 30 1-[(1-methylethyl)amino]-3-[2-(1H-pyrrol-1-yl)phenoxy]propan-2-ol;
- 1-(2-methoxyphenyl)-4-(4-succinimidobutyl)piperazine;
- 1-(2-methoxyphenyl)-4-(4-phthalimidobutyl)piperazine;
- 1-(1H-indol-4-yloxy)-3-[(1-methylethyl)amino]-2-propanol;
- (S)-1-(1H-indol-4-yloxy)-3-[(1-methylethyl)amino]-2-propanol;
- 35 N-[3-[3-(dimethylamino)ethoxy]-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-[1,1'-biphenyl]-4-carboxamide;
- 8-[(2,3-dihydro-1,4-benzodioxin-2-yl)methyl]-1-phenyl-1,3,8-triazaspiro [4,5]decan-4-one;
- (S)-N-tert-butyl-3-(4-(2-methoxyphenyl)-piperazin-1-yl)-2-phenylpropanamide;
- 8-[3-(4-fluorophenoxy)propyl]-1-phenyl-1,3,8-triazaspiro[4.5]-decan-4-one;
- 40 N-[2-[[3-(dimethylamino)propyl]thio]phenyl]-3-phenyl-2-propenamide;
- 4-(5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-methylpiperidine;
- 8-chloro-11-(1-piperazinyl)-5H-dibenzo[b,e][1,4]diazepine;
- 4-(4-fluorobenzoyl)-1-(4-phenylbutyl)-piperidine oxalate;
- 3-[2-[4-(4-fluorobenzoyl)-1-piperidinyl]ethyl]-2,4-[1H,3H]-quinazolinedione;
- 45 α -phenyl-1-(2-phenylethyl)-4-piperidinemethanol;
- metergoline phenylmethyl ester;
- 1,2,3,4,10,14b-hexahydro-2-methyldibenzo[c,f]pyrazino[1,2-a]azepine;
- 8-[5-(2,4-dimethoxy-5-(4-trifluoromethylphenylsulphonamido) phenyl-5-oxopentyl)]-1,3,8-triazaspiro[4.5]decane-2,4-dione;
- 50 N-(1-methyl-1H-indol-5-yl)-N'-3-pyridinyl-urea;
- N-(1-methyl-5-indolyl)-N'-(3-methyl-5-isothiazolyl)urea;
- (+)-cis-4,5,7a,8,9,10,11,11a-octahydro-7H-10-methylindolo[1,7-bc][2,6]-naphthyridine;
- 8-[4-(4-fluorophenyl)-4-oxobutyl]-1-phenyl-1,3,8-triazaspiro[4,5]decan-4-one;
- 55 6-chloro-2-[piperidinyl-4-thio]pyridine;
- 8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione;
- 1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine;
- 5-hydroxy-3-(1-methylpiperidin-4-yl)-1H-indole;
- 8-[4-[4-(2-pyrimidinyl)-1-piperazinyl]butyl]-8-azaspiro[4.5]decane-7,8-dione;
- 60 5-carboxamidotryptamine maleate;
- 7-trifluoromethyl-4-(4-methyl-1-piperazinyl)pyrrolo[1,2-a]-quinoxaline;

- 1,4-dihydro-3-(1,2,3,6-tetrahydro-4-pyridinyl)-5H-pyrrolo[3,2-b]pyridin-5-one;
 5-propoxy-3-(1,2,3,6-tetrahydro-4-pyridinyl)-1H-pyrrolo[3,2b]pyridine hydrochloride;
 3-[3-(2-dimethylaminoethyl)-1H-indol-5-yl]-N-(4-methoxybenzyl)acrylamide;
 65 8-hydroxy-2-dipropylaminotetralin hydrobromide;
 (2R)-(+)-8-hydroxy-2-(di-n-propylamino)tetralin;
 (RS)-trans-8-hydroxy-2-[N-n-propyl-N-(3'-iodo-2'-propenyl)amino]tetralin;
 2-[5-[3-(4-methylsulphonylamino)benzyl-1,2,4-oxadiazol-5-yl]-1H-indol-3-yl]ethanamine;
 70 8-[2-(1,4-benzodioxan-2-ylmethylamino)ethyl]-8-azaspiro[4.5]decane-7,9-dione;
 nonyloxytryptamineoxalate;
 5-methoxy-3-(1,2,5,6-tetrahydro-4-pyridinyl)-1H-indole;
 N-(3-trifluoromethylphenyl)piperazine;
 α -methyl-5-(2-thienylmethoxy)-1H-indole-3-ethanamine;
 75 1-(3-chlorophenyl)piperazine;
 α -methyl-5-hydroxytryptamine maleate; and
 6-chloro-2-(1-piperazinyl)pyrazine,
 or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof.

9. A composition comprising a cyclooxygenase-2 selective inhibitor selected from the group consisting of celecoxib, cimicoxib, deracoxib, valdecoxib, rofecoxib, lumiracoxib, etoricoxib, parecoxib, 2-(3,4-difluorophenyl)-4-(3-hydroxy-3-methylbutoxy)-5-[4-(methylsulfonyl)phenyl]-3(2H)-pyridazinone, and (S)-6,8-dichloro-2-(trifluoromethyl)-2H-
 5 1-benzopyran-3-carboxylic acid or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof; and a serotonin-modulating agent selected from the group consisting of:
 citalopram;
 fluoxetine;
 fluvoxamine;
 10 paroxetine;
 norfluoxetine;
 N-demethylsertraline;
 escitalopram oxalate;
 sertraline;
 15 3aS-2-[(S)-1-azabicyclo [2.2.2] oct-3-yl]-2,3,3a,4,5,6-hexahydro-1-oxo-1H-benz[de]isoquinoline hydrochloride;
 3-[4-(4-chlorophenyl)piperazin-1-yl]-1,1-diphenyl-2-propanol;

- 4-[3-[t-butylamino]-2-hydroxypropoxy]-1H-indole-2-carbonitrile;
3-[3-(dimethylamino)propyl]-4-hydroxy-N-[4-(4-pyridinyl) phenyl]benzamide;
20 N-[4-methoxy-3-(4-methyl-1-piperazinyl)phenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-1,1'-biphenyl-4-carboxamide hydrochloride;
1-[(1-methylethyl)amino]-3-[2-(1H-pyrrol-1-yl)phenoxy]propan-2-ol;
1-(2-methoxyphenyl)-4-(4-succinimidobutyl)piperazine;
1-(2-methoxyphenyl)-4-(4-phthalimidobutyl)piperazine;
25 1-(1H-indol-4-yloxy)-3-[(1-methylethyl)amino]-2-propanol;
(S)-1-(1H-indol-4-yloxy)-3-[(1-methylethyl)amino]-2-propanol;
N-[3-[3-(dimethylamino)ethoxy]-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-[1,1'-biphenyl]-4-carboxamide;
8-[(2,3-dihydro-1,4-benzodioxin-2-yl)methyl]-1-phenyl-1,3,8-triazaspiro [4,5]decan-
30 4-one;
(S)-N-tert-butyl-3-(4-(2-methoxyphenyl)-piperazin-1-yl)-2-phenylpropanamide;
8-[3-(4-fluorophenoxy)propyl]-1-phenyl-1,3,8-triazaspiro[4.5]-decan-4-one;
N-[2-[[3-(dimethylamino)propyl]thio]phenyl]-3-phenyl-2-propenamide;
4-(5H-dibenzo[a,d]cyclohepten-5-ylidene)-1-methylpiperidine;
35 8-chloro-11-(1-piperazinyl)-5H-dibenzo[b,e][1,4]diazepine;
4-(4-fluorobenzoyl)-1-(4-phenylbutyl)-piperidine oxalate;
3-[2-[4-(4-fluorobenzoyl)-1-piperidinyl]ethyl]-2,4[1H,3H]-quinazolinedione;
 α -phenyl-1-(2-phenylethyl)-4-piperidinemethanol;
metergoline phenylmethyl ester;
40 1,2,3,4,10,14b-hexahydro-2-methyldibenzo[c,f]pyrazino[1,2-a]azepine;
8-[5-(2,4-dimethoxy-5-(4-trifluoromethylphenylsulphonamido) phenyl-5-oxopentyl]-1,3,8-triazaspiro[4.5]decane-2,4-dione;
N-(1-methyl-1H-indol-5-yl)-N'-3-pyridinyl-urea;
N-(1-methyl-5-indolyl)-N'-(3-methyl-5-isothiazolyl)urea;
45 (+)-cis-4,5,7a,8,9,10,11,11a-octahydro-7H-10-methylindolo[1,7-bc][2,6]-naphthyridine;
8-[4-(4-fluorophenyl)-4-oxobutyl]-1-phenyl-1,3,8-triazaspiro[4,5]decan-4-one;
6-chloro-2-[piperidinyl-4-thio]pyridine;
8-[2-[4-(2-methoxyphenyl)-1-piperazinyl]ethyl]-8-azaspiro[4.5]decane-7,9-dione;
50 1-[3-(3,4-methylenedioxyphenoxy)propyl]-4-phenylpiperazine;
5-hydroxy-3-(1-methylpiperidin-4-yl)-1H-indole;
8-[4-[4-(2-pyrimidinyl)-1-piperazinyl]butyl]-8-azaspiro[4.5]decane-7,8-dione;

- 55 5-carboxamidotryptamine maleate;
 7-trifluoromethyl-4-(4-methyl-1-piperazinyl)pyrrolo[1,2-a]quinoxaline;
 1,4-dihydro-3-(1,2,3,6-tetrahydro-4-pyridinyl)-5H-pyrrolo[3,2-b]pyridin-5-one;
 5-propoxy-3-(1,2,3,6-tetrahydro-4-pyridinyl)-1H-pyrrolo[3,2b]pyridine hydrochloride;
 3-[3-(2-dimethylaminoethyl)-1H-indol-5-yl]-N-(4-methoxybenzyl)acrylamide;
 8-hydroxy-2-dipropylaminotetralin hydrobromide;
 (2R)-(+)-8-hydroxy-2-(di-n-propylamino)tetralin;
 60 (RS)-trans-8-hydroxy-2-[N-n-propyl-N-(3'-iodo-2'-propenyl)amino]tetralin;
 2-[5-[3-(-4-methylsulphonylamino)benzyl-1,2,4-oxadiazol-5-yl]-1H-indol-3-yl]ethanamine;
 8-[2-(1,4-benzodioxan-2-ylmethylamino)ethyl]-8-azaspiro[4.5]decane-7,9-dione;
 nonyloxytryptamineoxalate;
 65 5-methoxy-3-(1,2,5,6-tetrahydro-4-pyridinyl)-1H-indole;
 N-(3-trifluoromethylphenyl)piperazine;
 α -methyl-5-(2-thienylmethoxy)-1H-indole-3-ethanamine;
 1-(3-chlorophenyl)piperazine;
 α -methyl-5-hydroxytryptamine maleate; and
 70 6-chloro-2-(1-piperazinyl)pyrazine,
 or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof.

10. The composition of claim 9 wherein the cyclooxygenase-2 selective inhibitor is selected from the group consisting of celecoxib, cimicoxib, deracoxib, valdecoxib, rofecoxib, lumiracoxib, etoricoxib, and parecoxib; and the serotonin-modulating agent selected from the group consisting of citalopram, fluoxetine, fluvoxamine, paroxetine,
 5 norfluoxetine, N-demethylsertraline, escitalopram oxalate, sertraline, or an isomer, a pharmaceutically acceptable salt, ester, or prodrug thereof.

11. The composition of claim 10 wherein the cyclooxygenase-2 selective inhibitor is celecoxib and the serotonin-modulating agent is citalopram.

12. The composition of claim 10 wherein the cyclooxygenase-2 selective inhibitor is rofecoxib and the serotonin-modulating agent is citalopram.

13. The composition of claim 10 wherein the cyclooxygenase-2 selective inhibitor is valdecoxib and the serotonin-modulating agent is citalopram.

14. The method of any of claims 1-5 wherein the stroke is a hemorrhagic stroke.

15. The method of any of claims 1-5 wherein the stroke is an ischemic stroke.